

FIG.		PAGE
72.	RHIZOCRINUS LOFFOTENSIS, M. Sars. Once and a half the natural size. (No. 43, 1869)	451
73.	BATHYCRINUS GRACILIS (sp. n.). Twice the natural size. (No. 37, 1869)	453
74.	ARCHASTER BIFRONS (sp. n.). Oral aspect. Three-fourths the natural size. (No. 57, 1869)	455
75.	SOLASTER FURCIFER, Von Duben and Koren. Oral aspect. . Natural size. (No. 55, 1869)	456
76.	BUCCINOPSIS STRIATA, Jeffreys. Færoe Channel	464
77.	LATIRUS ALBUS, Jeffreys. Twice the natural size. Færoe Channel	464
78.	PLEURONECTIA LUCIDA, Jeffreys. Twice the natural size. <i>a</i> , from the Eastern Atlantic; <i>b</i> , from the Gulf of Mexico ...	465
79.	PECTEN HOSKYNSI, Forbes. Twice the natural size	465
80.	VENTRICULITES SIMPLEX, Toulmin Smith. Once and a half the natural size	483
81.	VENTRICULITES SIMPLEX, Toulmin Smith. Outer surface; four times the natural size	484
82.	VENTRICULITES SIMPLEX, Toulmin Smith. Section of the outer wall, showing the structure of the silicious network (x. 50)	485
83.	CÆLOSPHÆRA TUBIFEX (sp. n.). Slightly enlarged. Off the Coast of Portugal	485
84.	'CHOANITES.' In a flint from the white chalk	486

VIGNETTES.

THE FÆROE ISLANDS	xii
TINDHOLM	48
THORSHAVN	80
THE GOVERNOR'S HOUSE, THORSHAVN	132
LILLE DIMON	196
NOLSÖ, FROM THE HILLS ABOVE THORSHAVN	235
FUGLÖ, FROM THE EASTERN SHORE OF VIDERÖ	280
VAAÿ CHURCH IN SUDERÖ	328
THE GIANT AND THE HAG	406
BORDÖ, KUNÖ, AND KALSÖ, FROM THE HAMLET OF VIDERÖ	466
KUNÖ, FROM VAAÿ IN BORDÖ	501