

FIG.		PAGE
35.	<i>PHORMOSOMA HOPLACANTHA</i> (sp. n.). Southern Sea between Australia and New Zealand. Portion of the ventral surface of the test. Reduced one-third .....	148
36.	<i>UMBELLULARIA GREENLANDICA</i> , L. Natural size. (No. 7).....	150
37.	DIAGRAM SHOWING THE DIRECTION AND FORCE OF THE WIND (DIRECTION SOUTH-WEST BY COMPASS; FORCE = 7 BY BEAUFORT'S SCALE) AT MIDNIGHT, JANUARY 1ST, 1873 .....	158
38.	<i>POLIOPOGON AMADOU</i> (sp. n.). One-third of the natural size. (No. 3)....	171
39.	<i>ARCA</i> (sp.). (No. 5).....	174
40.	<i>LIMOPSIS</i> (sp.). (No. 5).....	174
41.	<i>LEDA</i> (sp.). (No. 5).....	175
42.	<i>WILLEMOESIA LEPTODACTYLA</i> , Von Willemoes-Suhm. Natural size. (No. 13). 183	
43.	<i>SOLARIUM</i> (sp.). Greatly enlarged. (No. 13).....	185
44.	<i>ANTENNARIUS MARMORATUS</i> . Natural size. From the surface.....	188
45.	<i>AVICULA</i> (sp.). Greatly enlarged. (No. 16).....	189
46.	<i>GLOBIGERINA BULLOIDES</i> , from the surface.....	202
47.	<i>ORBULINA UNIVERSA</i> , D'Orbigny. From the surface. Fifty times the natural size.....	205
48.	<i>PULVINULINA MENARDII</i> , D'Orbigny: <i>a</i> , the upper; <i>b</i> , the under surface. Thirty times the natural size. Dead shells from the bottom, at a depth of 1900 fathoms .....	208
49.	A "RHABDOSPHERE." From the surface. Five hundred times the natural size.....	210
50.	A "RHABDOSPHERE." From the surface. Two thousand times the natural size.....	211
51.	<i>CALCAROMMA CALCAREA</i> (sp. n.). With the pseudopodia contracted. From the surface. Two hundred times the natural size.....	220
52.	<i>DICTYOPODIUM</i> (sp. n.). From the surface. Two hundred times the natural size .....	221
53.	<i>XIPHACANTHA</i> (sp. n.). From the surface. One hundred times the natural size. The skeleton only.....	222
54.	<i>HALIOMMA</i> (sp. n.). From the surface. Two hundred times the natural size.....	223
55.	CURVES CONSTRUCTED FROM SERIAL SOUNDINGS BETWEEN TENERIFFE AND SOMBRERO.....	faces 226
56.	CURVES CONSTRUCTED FROM SERIAL AND BOTTOM SOUNDINGS. A, in the Bay of Biscay; B, off the coast of Portugal; and C and D, between Teneriffe and Sombrero .....	228
57.	DIAGRAM CONSTRUCTED FROM SERIAL SOUNDING No. 10.....	231
58.	DIAGRAM CONSTRUCTED FROM SERIAL SOUNDING No. 1.....	231
59.	<i>WILLEMOESIA CRUCIFER</i> , Von Willemoes-Suhm $\times$ 2. (No. 23).....	243
60.	<i>ASTACUS ZALECUS</i> , Von Willemoes-Suhm. (No. 23).....	245
61.	<i>CHETODERMA NITIDULUM</i> , Lovén. (No. 24).....	249
62.	<i>TROCHOCYATHUS CORONATUS</i> , Pourtales, $\times$ 2. (No. 24).....	250
63.	<i>DELTOCYATHUS AGASSIZII</i> , Pourtales, $\times$ 4. (No. 24).....	253
64.	<i>DELTOCYATHUS AGASSIZII</i> , Pourtales. Stellate variety. From a depth of 200 fathoms, near Bermudas.....	255
65.	<i>CRYPTOHELIA PUDICA</i> , Milne-Edwards. Twice the natural size. (No. 24)...	256
66.	<i>HYALONEMA TOXERES</i> (sp. n.). Upper surface, natural size. (No. 24)....	258