

- Lepidonotus wahlbergi*, Kinberg. Several specimens (Sea Point, between tide-marks); obtained at no other locality by the Challenger. Recorded from the Cape and Port Natal.
- Eunoa capensis*, n.sp. One specimen (Sea Point, between tide-marks); obtained at no other locality.
- Polynoë attenuata*, n.sp. One specimen (Sea Point, between tide-marks); obtained at no other locality.
- Eulalia capensis*, Schmarda. One specimen (Sea Point, between tide-marks); obtained at no other locality by the Challenger. Recorded from Table Bay.
- Notocirrus capensis*, n.sp. One specimen (Sea Point, between tide-marks); obtained at no other locality.
- Nematonereis* sp. (?). Fragment (Sea Point, beach).
- Eunice murrayi*, n.sp. One specimen (Simon's Bay, 18 fathoms); obtained at no other locality.
- Trophonia capensis*, n.sp. One specimen (Sea Point, between tide-marks); obtained at no other locality.
- Cirratulus capensis*, Schmarda. Several specimens (Sea Point, between tide-marks); obtained at no other locality by the Challenger. Recorded from the Cape.
- Nicomache capensis*, n.sp. One specimen (Sea Point, between tide-marks); obtained at no other locality.
- Sabellaria (Pallasia) capensis* (Schmarda). Several specimens (Sea Point, between tide-marks); obtained at no other locality by the Challenger. Recorded from the Cape.
- Schmardanella pterochæta* (Schmarda). Several specimens (Sea Point, between tide-marks); obtained at no other locality by the Challenger. Recorded from the Cape.
- Dasychone violacea* (Schmarda). Several specimens (Sea Point, between tide-marks); obtained at no other locality by the Challenger. Recorded from the Cape.

OSTRACODA (Brady, Zool. pt. 3).

- Pontocypris* (?) *subreniformis*, n.sp. (Simon's Bay, 15 to 20 fathoms); obtained also at Port Jackson, 2 to 10 fathoms.
- Macrocypris maculata*, Brady. (Simon's Bay, 15 to 20 fathoms); obtained also at Stations 145, 149, 162, and Amboina, 15 to 150 fathoms. Recorded from Australia, West Indies, and Turk's Island.