

Centrophorus crepidater, Boc. and Cap., 1902, Faroe Bank, 750 metres.

Centrophorus squamosus, Gmel., 1902, Faroe Bank, 390 to 750 metres (see Fig 256).




FIG. 256.

Centrophorus squamosus, Gmel. (After Jensen.)

Centrophorus calceus, Lowe, 1902, Faroe Bank, 750 metres.

Centrophorus coelolepis, Boc. and Cap., 1902, Faroe Bank, 750 metres.

Spinax niger, Bonap., 1902, Faroe Bank, 426 metres; 1910, Station 21.

Spinax (Etmopterus) princeps, Coll., 1902, Faroe-Shetland channel and Faroe Bank.

Centroscyllium fabricii (Reinh.), 1902, Faroe-Shetland channel and Faroe Bank.

RHINIDÆ

Rhina squatina, Duméril, 1910, Station 39.

Sub-Order—BATOIDEI

RAIIDÆ

Raia clavata, L. (thornback ray), 1902, Faroe Bank, 130 metres; 1910, Stations 1, 3, 13, 14, 20, 39 (see Fig. 257).




FIG. 257.

Raia clavata, L. (After Smitt.)

Raia punctata, Risso, 1910, Stations 37, 38, 39.

Raia microocellata, Montagu, 1910, Station 37.