

other was taken off the Congo. *Antennarius marmoratus* (see Fig. 471) presents in its shape some likeness to the Ceratiidæ and to *Lophius*, but is in fact a small surface form peculiar to the Sargasso Sea (see Plate V. Chapter X.), where the genus *Monacanthus*, belonging to the sub-order Plectognathi (see Fig. 472 and Plate V.), also occurs.

PELAGIC FISHES

TAKEN BY THE "MICHAEL SARS" IN 1910 IN THE ATLANTIC
NORTH OF LAT. 26° N.

Class—CYCLOSTOMATA

Order—PETROMYZONTES

PETROMYZONTIDÆ

Petromyzon marinus, L., 1910 (see Fig. 449).

Class—PISCES

Sub-Class—TELEOSTOMI

Order—TELEOSTEI

Sub-Order—MALACOPTERYGII

CLUPEIDÆ

Engraulis encrasicolus, Cuv., 1910, Station 36 (see Fig. 450).

Clupea alosa, L., 1910, Station 36.

Clupea pilchardus, Walb., 1910, Station 36 (see Fig. 451).

SALMONIDÆ

Opisthoproctus soleatus, Vaill., 1910, Stations 49, 52, 64 (see Fig. 72, p. 89).

Opisthoproctus grimaldii, Zugmayer, 1910, Stations 23, 49 (see Fig. 452).

STOMIATIDÆ

We give here Brauer's¹ classification of the Stomiidæ and Sternoptychidæ.

Stomias boa, Risso, 1902, Faroe-Shetland Channel; 1910, Stations 10, 19, 29, 34, 42, 51, 53, 56, 62, 63, 64, 67, 80, 82, 84, 87, 88, 90, 92, 94, 98, 101 (see Fig. 453).

Chauliodus sloanei, Bl. and Schn., 1910, Stations 10, 25, 29, 34, 35, 42, 45, 49, 50, 51, 52, 53, 56, 62, 63, 64, 80, 81, 82, 84, 88, 90 (see Fig. 454).

Photostomias guernei, Coll., 1910, Stations 34, 45, 49, 51, 53, 58, 81, 82 (see Fig. 455).

Eustomias obscurus, Vaill., 1910, Station 29.

¹ "Die Tiefsee Fische," *Wiss. Ergeb. Valdivia-Expedition*, Bd. 15, 1906.