

ENUMERATION OF THE PLANTS.

PHANEROGAMÆ.—ANGIOSPERMÆ.

DICOTYLEDONES.—POLYPETALÆ.

RANUNCULACEÆ.

Ranunculus muricatus, Linn.

Ranunculus muricatus, Linn., Sp. Pl., ed. 1, p. 555; Chapm., Fl. Southern U.S., p. 7.

BERMUDAS.—Introduced. Ditches and roadsides—*Lane; Jones; Lefroy; Moseley.*

An annual, common in Southern Europe and some parts of Asia, and now very widely spread as an introduced plant in America, Australia, &c.

Ranunculus parviflorus, Linn.

Ranunculus parviflorus, Linn., Sp. Pl., ed. 2, p. 780; Griseb., Fl. Brit. W. Ind., p. 2.

BERMUDAS.—Introduced. In similar situations to the last—*Lane; Jones; Rein; Lefroy; Moseley.*

This species is also an annual, and is now found in nearly all temperate and sub-tropical countries. They are both weeds of pasture rather than corn-land, and were early introduced into America, whence they may have been carried to the Bermudas.

PAPAVERACEÆ.

Argemone mexicana, Linn.

Argemone mexicana, Linn., Sp. Pl., ed. 1, p. 508; Griseb., Fl. Brit. W. Ind., p. 13; Desc., Fl. Ant., v. t. 380.

BERMUDAS.—Introduced. Very common—*Lane; Jones; Rein; Lefroy; Moseley.*

Annual. Almost universal in warm countries.

Papaver somniferum, Linn.

Papaver somniferum, Linn., Sp. Pl., ed. 1, p. 508; Gray, Manual, ed. 5, p. 59.

BERMUDAS.—Introduced. Occasionally as an escape from cultivation—*Lane; Jones; Lefroy.*

Cultivated and wild in many parts, especially of the Old World, but it is uncertain where it is really indigenous.

In Reade's collection there was a specimen of *Papaver dubium*, Linn.