

**Cakile æqualis**, L'Herit.

*Cakile æqualis*, L'Herit., in DC. Prodr., i. p. 185 ; Deless. Ic. Sel., ii. t. 57 ; Griseb., Fl. Brit. W. Ind., p. 14.

*Cakile maritima*, Scop., var. *æqualis*, Chapm., Fl. Southern U.S., p. 31.

BERMUDAS.—Indigenous. Common on the shores of the islands—*Lefroy* ; *Moseley* :

A maritime herb inhabiting the West Indies from Cuba to St Vincent and the Bahamas, and the Southern States of North America.

The European *Cakile maritima* is perhaps not specifically different. This is the scurvy-grass of the islanders, and most likely the plant named *Cochlearia officinalis* in Jones's list.

**Raphanus raphanistrum**, Linn.

*Raphanus raphanistrum*, Linn., Sp. Pl., ed. 1, p. 669 ; Gray, Manual, ed. 5, p. 75.

BERMUDAS.—Introduced. Without any note—*Moseley* ; *Reade*.

A weed of Europe, naturalised in many countries.

## CARYOPHYLLACEÆ.

**Cerastium vulgatum**, Linn.

*Cerastium vulgatum*, Linn., Sp. Pl., ed. 2, p. 627 ; Chapm., Fl. Southern U.S., p. 50 ; Gray, Manual, ed. 5, p. 93.

BERMUDAS.—Introduced. Cultivated ground—*Moseley* ; *Reade*.

A very common annual weed of cultivation in temperate and sub-tropical climates.

**Cerastium viscosum**, Linn.

*Cerastium viscosum*, Linn., Sp. Pl., ed. 1, p. 437 ; Griseb., Fl. Brit. W. Ind., p. 55 ; Chapm., Fl. Southern U.S., p. 50.

BERMUDAS.—Introduced. Without locality—*Lane*.

Also a very widely-dispersed annual weed.

**Stellaria media**, Cyr.

*Stellaria media*, Cyr., Char. Comm., p. 36 ; Griseb., Fl. Brit. W. Ind., p. 55 ; Chapm., Fl. Southern U.S., p. 50.

BERMUDAS.—Introduced. Cultivated ground—*Lane* ; *Rein* ; *Moseley*.

A very widely-dispersed annual in temperate regions.

**Arenaria serpyllifolia**, Linn.

*Arenaria serpyllifolia*, Linn., Sp. Pl., ed. 1, p. 463 ; Chapm., Fl. Southern U.S., p. 49.

BERMUDAS.—Introduced. A common weed—*Lefroy* ; *Reade*.

Widely dispersed in the Old World, and introduced in North America.