

**Melilotus parviflora, Desf.**

*Melilotus parviflora*, Desf., Fl. Atl., ii. p. 192; DC., Prodr., ii. p. 187.

BERMUDAS.—Introduced. Without locality—*Moseley*. Also in the British Museum, in Sloane's Herbarium, vol. xxxii. p. 83, collected by *Dickinson*.

An annual or biennial weed of cultivation from Europe.

**Vicia sativa, Linn.**

*Vicia sativa*, Linn., Sp. Pl., ed. 1, p. 736; Chapm., Fl. Southern U.S., p. 98; Gray, Manual, ed. 5, p. 138.

BERMUDAS.—Introduced. Common in pasture land—*Jones*.

A native of Europe and other parts of the Old World, and probably a relic of cultivation in these islands.

**Centrosema virginianum, Benth.**

*Centrosema virginianum*, Benth. in Mart. Fl. Bras., xv. 1, p. 132; Chapm., Fl. Southern U.S., p. 107.

BERMUDAS.—Indigenous? Without locality—*Rein*.

America, ranging from Maryland to South Brazil.

**Erythrina corallodendrum, Linn.**

*Erythrina corallodendrum*, Linn., Sp. Pl., ed. 1, p. 706; Griseb., Fl. Brit. W. Ind., p. 199; Desc., Fl. Ant., iv. t. 298.

*Erythrina speciosa*, Andr., Bot. Repos., t. 443.

BERMUDAS.—Introduced. Introduced into the Bermudas from St Thomas about fifty years ago—*Darrell*.

West Indies to Brazil.

**Canavalia obtusifolia, DC.**

*Canavalia obtusifolia*, DC., Prodr., ii. p. 404; Griseb., Fl. Brit. W. Ind., p. 197; Desc., Fl. Ant., viii. t. 559; Chapm., Fl. Southern U.S., p. 109.

*Canavalia rosea* et *Canavalia miniata*, DC., loc. cit.

BERMUDAS.—Indigenous. Common on the sea-coast—*Jones*; *Moseley*; *Lefroy*.

This is a very common sea-shore plant in nearly all warm countries, including many remote islands. It appears to be one of those *Leguminosæ* whose seeds germinate after long immersion in sea-water. It bears the local name of "Bay-bean."

**Vigna luteola, Benth.**

*Vigna luteola*, Benth. in Mart. Fl. Bras., xv. 1, p. 194, t. 50, fig. 2; Griseb., Fl. Brit. W. Ind., p. 195.

*Vigna glabra*, Savi; Chapm., Fl. Southern U.S., p. 106.

BERMUDAS.—Indigenous? Without locality—*Lefroy*.