

A common American plant in brackish marshes on the sea-shores, from Carolina to Buenos Ayres and Chili; also in Africa, Australia, &c.

*Cajanus indicus*, Spreng.

*Cajanus indicus*, Spreng., Syst. Veg., iii. p. 248; Griseb., Fl. Brit. W. Ind., p. 191.

BERMUDAS.—Introduced. Common on David Island—*Jones*; *Lefroy*.

Believed to be a native of the Old World; but it is now wild in most warm countries, having spread from cultivation.

*Sophora tomentosa*, Linn.

*Sophora tomentosa*, Linn., Sp. Pl., ed. 1, p. 373; Griseb., Fl. Brit. W. Ind., p. 203; Chapm., Fl. Southern U.S., p. 113; Desc., Fl. Ant., vii. t. 475.

BERMUDAS.—Indigenous. Without locality—*Lefroy's list*; Boaz Island—*Reade*.

There is also a specimen in the Herbarium of the British Museum labelled "Bermuda," but without the collector's name.

This tree occurs on nearly all tropical sea-shores, and extends beyond the tropics in some parts of the world.

CÆSALPINIÆ.

*Cæsalpinia bonducella*, Fleming.

*Cæsalpinia bonducella*, Fleming in Asiat. Res., xi. p. 159.

*Guilandina bonducella*, Linn., Sp. Pl., ed. 2, p. 545; Griseb., Fl. Brit. W. Ind., p. 204; Desc., Fl. Ant., ii. t. 90.

BERMUDAS.—Introduced? Walsingham—*Lefroy*.

A cosmopolitan shrub in warm climates, flourishing equally well on the sea-shore and inland, ascending to an altitude of 2500 feet in the Himalayas.

*Cassia occidentalis*, Linn.

*Cassia occidentalis*, Linn., Sp. Pl., ed. 1, p. 377; Griseb., Fl. Brit. W. Ind., p. 209; Chapm., Fl. Southern U.S., p. 114; Desc., Fl. Ant., ii. t. 135.

BERMUDAS.—Introduced. Common—*Lefroy*.

An annual species, generally diffused in warm countries, except Australasia.

*Cassia bicapsularis*, Linn.

*Cassia bicapsularis*, Linn., Sp. Pl., ed. 1, p. 376; Griseb., Fl. Brit. W. Ind., 207; Jacq., Fragm., t. 58.

BERMUDAS.—Introduced. Without localities—*Moseley*; *Lefroy*.

A shrubby species, having a very wide range in the warmer parts of America and the West Indian islands.