

Psidium pomiferum et Psidium pyriferum, Linn.

Psidium pomiferum et Psidium pyriferum, Linn., Sp. Pl., ed. 2, p. 672.

Psidium guava, Radd. ex Griseb. Fl. Brit. W. Ind., p. 241.

BERMUDAS.—Introduced. Marsh near Devonshire—*Lane*.

The Guava, a native of Tropical America, is naturalised in all warm countries.

From a letter to Sir J. H. Lefroy, signed W. Cox, it would appear that for many years past nobody has troubled to cultivate it in the Bermudas.

Punica granatum, Linn.

Punica granatum, Linn., Sp. Pl., ed. 1, p. 472; Griseb., Fl. Brit. W. Ind., p. 242.

BERMUDAS.—Introduced—*Lane*; *Lefroy*.

Mediterranean region, and eastward to North-Western India, where it is believed to be indigenous.

ONAGRARIEÆ.

Oenothera rosea, Ait.

Oenothera rosea, Ait., Hort. Kew, ed. 1, ii. p. 3; Bot. Mag., t. 347; Griseb., Fl. Brit. W. Ind., p. 273.

BERMUDAS.—Introduced. Without locality—*Moseley*; *Reade*.

Texas to New Grenada, and in the West Indies. Naturalised in many other countries.

Oenothera sinuata, Michx.

Oenothera sinuata, Michx., Fl. Bor.-Am., i. p. 224; Chapm., Fl. Southern U.S., p. 138.

Oenothera humifusa, Nutt., Gen. N. Am. Pl., i. p. 245.

BERMUDAS.—Introduced. Roadsides, &c.—*Rein*; *Lefroy*; *Reade*.

Southern States of North America and Mexico.

Gaura coccinea, Nutt.

Gaura coccinea, Nutt. in Torr. and Gray, Fl. N. Am., i. p. 73; Hemsl. in Godm. and Salv., Biol. Centr. Am., i. p. 466.

BERMUDAS.—Introduced. Without locality—*Rein*.

Widely dispersed in Western North America, including Mexico.

Ludwigia repens, Swartz.

Ludwigia repens, Swartz, Fl. Ind. Occ., i. p. 273; Ic. Pl. Ind. Occ., t. 8.

Isardia repens, DC., Prodr., iii. p. 60; Griseb., Fl. Brit. W. Ind., p. 271.

Ludwigia natans, Ell; Chapm., Fl. Southern U.S., p. 142, fide Griseb., Fl. Brit. W. Ind., p. 271.

BERMUDAS.—Indigenous. Abundant in peat marshes—*Moseley*; without locality—*Rein*.

South Carolina to Texas and the West Indies.