

Chiococca racemosa, Linn.

Chiococca racemosa, Linn., Syst. Nat., ii. p. 917; Jacq., Sel. Stirp. Am., p. 68; Griseb., Fl. Brit. W. Ind., p. 336; Chapm., Fl. Southern U.S., p. 177.

BERMUDAS.—Indigenous. Walsingham—*Lane*; abundant in places—*Lefroy*; *Moseley*; roadside near Tucker's Town—*Jones*; in the bays associated with *Dodonæa viscosa*, &c.—*Rein.*

Florida and Mexico southward to Peru and Brazil.

The "Blolly" of the islanders.

Coffea arabica, Linn.

Coffea arabica, Linn., Sp. Pl., ed. 1, p. 172; Griseb., Fl. Brit. W. Ind., p. 338.

BERMUDAS.—Introduced. Grows wild about Walsingham and different parts of the islands—*Jones*; confined to the Cave district—*Lefroy*; without locality—*Rein.*

Eastern Tropical Africa; also naturalised in the West Indies.

Morinda royoc, Linn.

Morinda royoc, Linn., Sp. Pl., ed. 1, p. 176; Griseb., Fl. Brit. W. Ind., p. 347; Chapm., Fl. Southern U.S., p. 177.

BERMUDAS.—Indigenous. Walsingham—*Lane*; without locality—*Lefroy*.

Florida, Cuba, Jamaica, Honduras, and Panama, usually growing in maritime districts.

Psychotria undata, Jacq.

Psychotria undata, Jacq., Hort. Schœnb., iii. p. 5, t. 260; Griseb., Fl. Brit. W. Ind., p. 342; Chapm., Fl. Southern U.S., p. 177.

BERMUDAS.—Indigenous. Walsingham—*Lefroy*; without locality—*Moseley*.

Florida, Bahamas, and Jamaica.

Spermacoce tenuior, Lam.

Spermacoce tenuior, Lam. ex Griseb. Fl. Brit. W. Ind., p. 349; Chapm., Fl. Southern U.S., p. 175.

BERMUDAS.—Introduced? Without localities or memoranda—*Lefroy*; *Reade*.

An annual herb, inhabiting Florida and Mexico and the country southward to Peru and Brazil.

Vaillantia hispida, Linn.

Vaillantia hispida, Linn., Sp. Pl., ed. 2, p. 1490; DC., Prodr., iv. p. 614.

BERMUDAS.—Introduced. Walls and roadsides—*Moseley*; *Rein.*

Introduced from Europe.

Lane's list includes "*Vaillantia muralis*," Linn., probably in error for this species.