

Relbunium hypocarpium, Hemsl.

Relbunium hypocarpium, Hemsl. in Salv. and Godm. Biolog. Centr. Am. Bot., ii. p. 63.

Rubia hypocarpia, DC., Prodr., iv. p. 591.

Galium hypocarpium, Endl. ex Griseb. Fl. Brit. W. Ind., p. 351.

BERMUDAS.—Introduced? Without special locality—*Moseley*; *Lefroy*.

Generally dispersed in the West Indies and Tropical South America.

Galium uniflorum, Michx.

Galium uniflorum, Michx., Fl. Bor.-Am., i. p. 79; Chapm., Fl. Southern U.S., p. 174.

BERMUDAS.—Indigenous. In various localities—*Munro*; *Lefroy*; *Reade*.

South Carolina to Florida.

Galium bermudense, Linn.

Galium bermudense, Linn., Sp. Pl., ed. 1, p. 105.

Rubia tetraphylla, glabra, latiore folio, bermudensis, seminibus binis atropurpureis.—Pluk., Alm., p. 324, et Phytogr., t. 248. "A Do. Petiver habuimus."

It is uncertain what species Plukenet's plant is. Pursh (Fl. Am. Sept., i. p. 104) regards *Galium punctulosum*, Michx. (Fl. Bor.-Am., i. p. 8), and *Galium purpureum*, Walt. (Fl. Carol., p. 87), as the same. Chapman (Fl. Southern U.S., p. 174) cites *Galium bermudianum*, Ell. (Bot. Carol., i. p. 196), under *Galium pilosum*, Ait.

VALERIANEÆ.

Centranthus macrosiphon, Boiss.

Centranthus macrosiphon, Boiss., Diag. Pl. Nov., iii. p. 57; Walp., Rep., vi. p. 80; Lem., Jard. Fleur., iii. t. 273.

BERMUDAS.—Introduced. Without locality—*Rein*.

A native of Western Europe, probably imported into the islands for its beauty.

Valerianella olitoria, Mœnch.

Valerianella olitoria, Mœnch., Meth. Pl., p. 493; DC., Prodr., iv. p. 625; Gray, Manual, ed. 5, p. 214 (*Fedia*).

Fedia radiata, Michx. ? Chapm., Fl. Southern U.S., p. 184.

BERMUDAS.—Introduced. St David Island—*Lefroy*.

Mediterranean region, and naturalised in North America and other countries where it is cultivated.