

Erigeron philadelphicus, Linn.

Erigeron philadelphicus, Linn., Sp. Pl., ed. 1, p. 863 ; Chapm., Fl. Southern U.S., p. 206.

BERMUDAS.—Introduced? A common weed—*Lane*; *Jones*; *Rein*; *Lefroy*.

Canada to Florida.

Erigeron tenuis, Torr. and Gray.

Erigeron tenuis, Torr. and Gray, Fl. N. Am., ii. p. 175.

Erigeron quercifolius, Nutt. in DC. Prodr., v. p. 285 sed certe non Lam., ex Torr. and Gray.

BERMUDAS.—Introduced. Without locality—*Moseley*. Not included in any of the other collections, and probably only of casual occurrence in the islands.

Louisiana ; Texas.

Baccharis glomeruliflora, Pers.

Baccharis glomeruliflora, Pers., Ench., ii. p. 423 ; Chapm., Fl. Southern U.S., p. 218.

Baccharis sessiliflora, Michx., Fl. Bor.-Am., ii. p. 125.

BERMUDAS.—Indigenous. Common in marshes—*Farlow*; *Lane*; *Moseley*; *Lefroy*.

North Carolina to Florida.

This is the "Dogwood" of the islanders; and the *Baccharis*, n. sp.? of Jones's list, and the *Baccharis heterophylla*, Linn., of Rein's list, is probably the same species, for there appears to be only one in the islands. "*Baccharis halimifolia*, Linn.," of Lane's list, is also doubtless the same species.

Pluchea odorata, Cass.

Pluchea odorata, Cass., Dict., xlii. p. 3 ; DC., Prodr., v. p. 452 ; Griseb., Fl. Brit. W. Ind., p. 366 ; Desc., Fl. Ant., iii. t. 217.

BERMUDAS.—Indigenous? Pembroke Marsh—*Lefroy*.

Florida and Mexico to New Grenada, and throughout the West Indies.

Lefroy states that this grows eight feet high, with a stem as thick as a man's wrist.

Pluchea purpurascens, DC.

Pluchea purpurascens, DC., Prodr. v. p. 452 ; Griseb. Fl. Brit. W. Ind., p. 367 ; Chapm., Fl. Southern U.S., p. 218.

BERMUDAS.—Indigenous. Shelly Bay and Warwick—*Lefroy*.

Florida and Mexico to New Grenada, and throughout the West Indies.

Pluchea camphorata, DC.

Pluchea camphorata, DC., Prodr., v. p. 452 ; Chapm., Fl. Southern U.S., p. 218 ; Gray, Manual, ed. 5, p. 247.

BERMUDAS.—Indigenous. Marshes—*Moseley*.