

An inhabitant of salt marshes in eastern North America, ranging from North Carolina to Florida.

Polymnia uvedalia, Linn.

Polymnia uvedalia, Linn., Sp. Pl., ed. 2, p. 1303; Chapm., Fl. Southern U.S., p. 219; Gray, Manual, ed. 5, p. 248.

BERMUDAS.—Introduced. Escaped from cultivation—*Lane*; without locality—*Rein*; *Reade*.

New York to Florida.

Parthenium hysterophorus, Linn.

Parthenium hysterophorus, Linn., Sp. Pl., ed. 1, p. 988; Griseb., Fl. Brit. W. Ind., p. 369; Chapm., Fl. Southern U.S., p. 222.

BERMUDAS.—Introduced? Common—*Lefroy*; *Rein*; *Moseley*; *Reade*.

Florida and Mexico to Patagonia, chiefly in waste places and on roadsides.

Ambrosia artemisiæfolia, Linn.

Ambrosia artemisiæfolia, Linn., Sp. Pl., ed. 1, p. 988; Chapm., Fl. Southern U.S., p. 223; Griseb., Fl. Brit. W. Ind., p. 370.

BERMUDAS.—Introduced. Roadsides—*Lane*; *Lefroy*.

Canada to Brazil, especially in cultivated ground and waste places.

The "*Ambrosia heterophylla*, DC.," of Jones's list is perhaps the same species.

Eclipta erecta, Linn.

Eclipta erecta, Linn. ex DC. Prodr., v. p. 490; Chapm., Fl. Southern U.S., p. 224; Griseb., Fl. Brit. W. Ind., p. 370, sub *Eclipta alba*.

BERMUDAS.—Indigenous? Marshes—*Moseley*; abundant in wet ditches—*Lefroy*.

Pennsylvania to Uruguay, and widely diffused in the Old World.

Borrichia arborescens, DC. (Plate II.)

Borrichia arborescens, DC., Prodr., v. p. 489; Griseb., Fl. Brit. W. Ind., p. 371; Chapm., Fl. Southern U.S., p. 224.

Buphthalmum arborescens, Linn., Sp. Pl., ed. 2, p. 1273.

Chrysanthemum bermudense, folio rigido-viridi, Moris., Hist. Pl., pars iii. p. 25.

Borrichia argentea, DC., Prodr., v. p. 489.

BERMUDAS.—Indigenous. Abundant along the whole coast-line—*Rein*; *Jones*; *Moseley*; *Lefroy*; Boaz Island—*Reade*.

The Keys of Florida, and through the West Indies to Peru.