

**Dichondra repens, Forst.**

*Dichondra repens*, Forst., Char. Gener., p. 40, t. 20; Griseb., Fl. Brit. W. Ind., p. 476; Chapm. Fl. Southern U.S., p. 346.

BERMUDAS.—Indigenous. Marshes—*Moseley*; pastures—*Lane*; without locality—*Rein*.

Generally diffused in Temperate and Tropical America, Tropical Asia, Southern Africa, Australia, New Zealand, and the Pacific Islands.

## SOLANACEÆ.

**Solanum aculeatissimum, Jacq.**

*Solanum aculeatissimum*, Jacq., Ic. Pl. Rar., t. 41; Griseb., Fl. Brit. W. Ind., p. 442; Chapm., Fl. Southern U.S., p. 349.

BERMUDAS.—Introduced? Without locality—*Rein*.

A widely spread species in Tropical and Sub-tropical America.

Mr Moseley collected an imperfect specimen of a *Solanum*, which may be this, or *Solanum mammosum*, Linn.; and Sir J. H. Lefroy records *Solanum torvum*, Sw., in his list, on the authority of the late General Munro. The latter is also in Jones's list.

**Solanum nigrum, Linn.**

*Solanum nigrum*, Linn., Sp. Pl., ed. 1, p. 186; Chapm., Fl. Southern U.S., p. 348.

*Solanum nodiflorum*, Jacq., Ic. Pl. Rar., t. 326; Griseb., Fl. Brit. W. Ind., p. 437.

BERMUDAS.—Introduced. Waste places—*Lane*; *Rein*.

This is now generally diffused in temperate and tropical regions throughout the world.

**Nicandra physaloides, Gærtn.**

*Nicandra physaloides*, Gærtn., Fruct., ii. p. 237, t. 131; Chapm., Fl. Southern U.S., p. 351.

BERMUDAS.—Introduced. Ireland Island—*Lane*.

Peru, and naturalised in most warm countries.

**Physalis lanceolata, Michx.**

*Physalis lanceolata*, Michx., Fl. Bor.-Am., i. p. 149; Chapm., Fl. Southern U.S., p. 350.

BERMUDAS.—Introduced. Walsingham—*Lane*.

Florida to North Carolina, in dry sandy soil and near the sea.

[*Physalis peruviana*, Linn., a commonly cultivated and widely spread species, is included in Jones's list.]