

**Linaria vulgaris, Mill.**

*Linaria vulgaris*, Mill., Dict.; DC., Prodr., x. p. 273; Chapm., Fl. Southern U.S., p. 290; Griseb., Fl. Brit. W. Ind., p. 431.

BERMUDAS.—Introduced. Common in cultivated ground—*Jones; Reade*.  
Europe and Northern Asia; introduced in North America and elsewhere.

**Pentstemon pubescens, Solander.**

*Pentstemon pubescens*, Solander in Ait. Hort. Kew, ed. 1, iii. p. 360; Chapm., Fl. Southern U.S., p. 290.

BERMUDAS.—Introduced. Without locality—*Rein*.  
A native of eastern North America, ranging from Canada to Florida and Texas.

**Herpestis monniera, H. B. K.**

*Herpestis monniera*, H. B. K., Nov. Gen. et Sp., ii. p. 366; Griseb., Fl. Brit. W. Ind., p. 430; Chapm., Fl. Southern U.S., p. 292.

BERMUDAS.—Indigenous. Somerset Island—*Lefroy*; without locality—*Rein; Reade*.  
Nearly all tropical and sub-tropical countries; inhabiting marshy and maritime places.

**Capraria biflora, Linn.**

*Capraria biflora*, Linn., Sp. Pl., ed. 1, p. 628; Griseb., Fl. Brit. W. Ind., p. 427; Chapm., Fl. Southern U.S., p. 296.

BERMUDAS.—Introduced. Paget parish—*Lefroy*; Hamilton—*Lane*.  
Florida and Mexico to Peru and Brazil.

Locally known as "Tea-plant." "Introduced into the islands about fifty years ago by Dr Hinson."—J. H. Darrell in a communication to Sir J. H. Lefroy.

**Veronica agrestis, Linn.**

*Veronica agrestis*, Linn., Sp. Pl., ed. 1, p. 13; Chapm., Fl. Southern U.S., p. 296; Gray, Manual, ed. 5, p. 333.

BERMUDAS.—Introduced. *Lane; Lefroy; Reade*.

North temperate zone in the Old World, and naturalised in many other parts of the world.

**Veronica arvensis, Linn.**

*Veronica arvensis*, Linn., Sp. Pl., ed. 1, p. 13; Chapm., Fl. Southern U.S., p. 296.

BERMUDAS.—Introduced. *Moseley; Lefroy*.

North temperate zone in the Old World, and introduced into other countries.