

Verbena urticæfolia, Linn.

Verbena urticæfolia, Linn., Sp. Pl., ed. 1, p. 20; Griseb., Fl. Brit. W. Ind., p. 493; Chapm., Fl. Southern U.S., p. 307.

BERMUDAS.—Indigenous? Without locality—*Moseley*. Also in the Sloane Herbarium, clix. p. 47 (“a strong emetic”).

Canada to Florida, Mexico, and the West Indies.

Lefroy records *Verbena officinalis*, Linn., in his list, a plant not unlikely to occur.

[*Citharexylum quadrangulare*, Jacq., is in some collections, whether wild or not has not been ascertained.]

Duranta plumieri, Jacq.

Duranta plumieri, Jacq., Sel. Stirp. Am., p. 186, t. 176, fig. 76; Griseb., Fl. Brit. W. Ind., p. 498; Chapm., Fl. Southern U.S., p. 309.

BERMUDAS.—Indigenous? Quite common—*Lefroy*; *Moseley*; *Reade*.

Florida and Mexico to Brazil; also in Tropical Africa.

[In the Banksian collection at the British Museum, there is a flowering specimen of a *Petræa* (*Petræa volubilis*?) from “Bermudas.”]

Avicennia nitida, Jacq.

Avicennia nitida, Jacq., Sel. Stirp. Am., p. 177, t. 112, fig. 1; Griseb., Fl. Brit. W. Ind., p. 502; Gray, Synop. Fl. N. Am., ii. p. 341.

Avicennia tomentosa, Nutt. ? and *Avicennia tomentosa*, Jacq. in Chapm., Fl. Southern U.S., p. 310.

BERMUDAS.—Indigenous. Everywhere in salt marshes associated with *Rhizophora*—*Lane*; *Rein*; *Jones*; *Moseley*; *Lefroy*.

Florida, West Indies, and Mexico to Brazil; also in Tropical Africa.

Locally known as “Black Mangrove.”

LABIATÆ.**Mentha rotundifolia, Linn.**

Mentha rotundifolia, Linn., Sp. Pl., ed. 2, p. 805; Chapm., Fl. Southern U.S., p. 312.

BERMUDAS.—Introduced. Wild by roadsides—*Lefroy*; *Reade*.

Europe, North Africa, and Temperate Asia, though frequently occurring only as a colonist.

Pycnanthemum muticum, Pers.

Pycnanthemum muticum, Pers., Ench., ii. p. 128; Chapm., Fl. Southern U.S., p. 315.

Brachystemum muticum, Michx., Fl. Bor.-Am., ii. p. 6, t. 32.

BERMUDAS.—Introduced. Without locality—*Rein*.

Maine to Ohio and Florida, usually growing in dry or sandy soil.