

Calamintha officinalis, Mœnch.

Calamintha officinalis, Mœnch, Meth. Pl., p. 409 ; DC., Prodr., xii. p. 228.
Melissa calamintha, Linn., Sp. Pl., ed. 1, p. 593.

BERMUDAS.—Introduced. Without locality—*Rein.*

North temperate regions of the Old World, formerly commonly cultivated.

Salvia coccinea, Linn.

Salvia coccinea, Linn., fl., Sp. Pl. Suppl., p. 88 ; Griseb., Fl. Brit. W. Ind., p. 490 ; Gray, Synop. Fl. N. Am., ii. p. 368.

BERMUDAS.—Introduced. Common—*Lane* ; *Rein* ; *Moseley*.

Carolina to Florida (" but probably introduced "—*A. Gray*), West Indies, Mexico, and South America.

Salvia occidentalis, Swartz.

Salvia occidentalis, Swartz, Fl. Ind. Occ., i. p. 43 ; Griseb., Fl. Brit. W. Ind., p. 490 ; Gray, Synop. Fl. N. Am., ii. p. 370.

BERMUDAS.—Introduced ? Walsingham—*Lane* ; without locality—*Moseley*.

Florida, West Indies, and Mexico to Peru, abounding in waste places and as a weed of cultivation.

Salvia serotina, Linn.

Salvia serotina, Linn., Mant., p. 25 ; Griseb., Fl. Brit. W. Ind., p. 490 ; Chapm., Fl. Southern U.S., p. 319.

BERMUDAS.—Introduced ? Roadsides, &c.—*Lane* ; *Rein* ; *Moseley*.

South Florida, and throughout the West Indies.

Marrubium vulgare, Linn.

Marrubium vulgare, Linn., Sp. Pl., ed. 1, p. 583 ; Gray, Synop. Fl. N. Am., ii. p. 384.

BERMUDAS.—Introduced. *Rein.*

North temperate regions of the Old World, and introduced into North America.

Sideritis romana, Linn.

Sideritis romana, Linn., Sp. Pl., ed. 1, p. 575 ; DC., Prodr., xii. p. 445 ; Sibthorp, Fl. Græc., t. 552.

BERMUDAS.—Introduced. *Rein.*

An inhabitant of the Mediterranean region, and probably only of casual occurrence in the islands.

Leonurus sibiricus, Linn.

Leonurus sibiricus, Linn., Sp. Pl., ed. 1, p. 584 ; Griseb., Fl. Brit. W. Ind., p. 491 ; Gray, Synop. Fl. N. Am., ii. p. 385.

BERMUDAS.—Introduced. Roadsides—*Moseley* ; *Lefroy*.

Europe and Siberia, and naturalised in many other countries.