

***Urtica chamædryoides*, Pursh.**

Urtica chamædryoides, Pursh., Fl. Am. Sept., i. p. 113; DC., Prodr., xvi. 1, p. 42 (varietates);
Chapm., Fl. Southern U.S., p. 412.

BERMUDAS.—Indigenous? Without locality—*Lefroy*.

Georgia; Texas; Mexico.

***Urtica dioica*, Linn.**

Urtica dioica, Linn., Sp. Pl., ed. 1, p. 984; Chapm., Fl. Southern U.S., p. 412; Gray, Manual,
ed. 5, p. 444.

BERMUDAS.—Introduced. *Rein; Reade*.

A domestic plant of the Old World, now wild in most temperate and sub-tropical regions.

***Urtica urens*, Linn.**

Urtica urens, Linn., Sp. Pl., ed. 1, p. 984; Chapm., Fl. Southern U.S., p. 412.

BERMUDAS.—Introduced. *Rein; Moseley*.

Old World, and now nearly as widely spread as the last.

***Boehmeria cylindrica*, Willd.**

Boehmeria cylindrica, Willd., Sp. Pl., iv. p. 340; Griseb., Fl. Brit. W. Ind., p. 160; Chapm., Fl.
Southern U.S., p. 414.

BERMUDAS.—Indigenous. Marshes, common—*Moseley; Lefroy*.

Canada to Brazil and Uruguay; exceedingly common in many parts in swampy and shady situations.

***Parietaria officinalis*, Linn.**

Parietaria officinalis, Linn., Sp. Pl., ed. 1, p. 1052; DC., Prodr., xvi. 1, p. 235⁴².

BERMUDAS.—Introduced. Common—*Moseley; Reade*.

Introduced from Europe.

***Parietaria debilis*, Forst.**

Parietaria debilis, Forst., Fl. Ins. Austr. Prodr., p. 73; Chapm., Fl. Southern U.S., p. 414.

BERMUDAS.—Indigenous. Walls in St. George's—*Lefroy*; without locality—*Rein*.

Very widely diffused in temperate and warm regions.

***Parietaria pennsylvanica*, Mühlenb.**

Parietaria pennsylvanica, Mühlenb., Cat. Pl. Am. ex Wedd. in DC. Prodr., xvi. 1, p. 235⁴⁷;
Chapm., Fl. Southern U.S., p. 413.

BERMUDAS.—Indigenous. North shore—*Lefroy*; without locality—*Moseley*.

Vermont to Wisconsin, and southward to Florida and Mexico.