

Heleocharis capitata, R. Br.

Heleocharis capitata, R. Br., Prodr. Fl. N. Holl., p. 225; Chapm., Fl. Southern U.S., p. 518 (*Eleocharis*).
Scirpus capitatus, Willd., Sp. Pl., i. p. 294; Griseb., Fl. Brit. W. Ind., p. 570.

BERMUDAS.—Indigenous. Marshes—*Moseley*.

Tropical and Sub-tropical America, Asia, Africa, and Australia.

Heleocharis melanocarpa, Torr.

Heleocharis melanocarpa, Torr. in Ann. Lyc. Nat. Hist. N. York, iii. p. 311; Chapm., Fl. Southern U.S., p. 516 (*Eleocharis*).

BERMUDAS.—Indigenous. *Rein*.

North America, on the coast from Massachusetts to Florida.

Heleocharis plantaginea, R. Br.

Heleocharis plantaginea, R. Br., Prodr. Fl. N. Holl., p. 224, *in nota* (*Eleocharis*).

Scirpus plantagineus, Linn. ex Griseb., Fl. Brit. W. Ind., p. 571.

Eleocharis equisetoides, Torr. in Ann. Lyc. Nat. Hist. N. York, iii. p. 296; Chapm., Fl. Southern U.S., p. 514.

BERMUDAS.—Indigenous. Marshes—*Lefroy*.

Florida and Louisiana to Brazil.

Dichromena leucocephala, Michx.

Dichromena leucocephala, Michx., Fl. Bor.-Am., i. p. 37; Chapm., Fl. Southern U.S., p. 530; Gray, Manual, ed. 5, p. 567.

Rhynchospora stellata, Griseb., Fl. Brit. W. Ind., p. 576.

BERMUDAS.—Indigenous. Marshes, &c.—*Moseley*; *Lefroy*; *Rein*; *Reade*.

New Jersey to Florida and the West Indies.

Dichromena pura, Nees.

Dichromena pura, Nees in Linnæa, ix. p. 291; Kunth, Enum., ii. p. 283 (*nomen tantum*).

Rhynchospora pura, Griseb., Fl. Brit. W. Ind., p. 577.

BERMUDAS.—Indigenous. *Rein*.

There are no Bermudan specimens of this species in the Kew Herbarium.

West Indies, Venezuela, and Guiana.

Scirpus lacustris, Linn.

Scirpus lacustris, Linn., Sp. Pl., ed. 1, p. 48; Chapm., Fl. Southern U.S., p. 520; Boeckl. in Linnæa, xxxvi. p. 712.

Scirpus validus, Vahl, Enum., ii. p. 268; Griseb., Fl. Brit. W. Ind., p. 571; Gray, Manual, ed. 5, p. 562.

BERMUDAS.—Indigenous. Salt marshes—*Moseley*; *Reade*.

Generally diffused in temperate and sub-tropical countries.