

**Panicum capillare**, Linn.

*Panicum capillare*, Linn., Sp. Pl., ed. 1, p. 58; Chapm., Fl. Southern U.S., p. 574; Gray, Manual, ed. 5, p. 647.

BERMUDAS.—Introduced. *Rein.*

Eastern North America, growing in sandy soil and on the coast, from New Jersey southward.

**Panicum paspaloides**, Pers.

*Panicum paspaloides*, Pers., Ench., i. p. 81; Griseb., Fl. Brit. W. Ind., p. 545; Kunth, Enum., i. 1, p. 77.

BERMUDAS.—Introduced. *Moseley.*

Widely distributed in warm countries.

**Panicum horizontalis**, Mey.

*Panicum horizontalis*, Mey., Esseq., p. 54; Kunth, Enum., i. 1, p. 81.

*Digitaria setigera*, Roth, Nov. Sp., p. 37; Griseb., Fl. Brit. W. Ind., p. 544.

BERMUDAS.—Introduced. *Rein.*

West Indies and other warm countries.

**Panicum virgatum**, Linn.

*Panicum virgatum*, Linn., Sp. Pl., p. 59; Chapm., Fl. Southern U.S., p. 573; Gray, Manual, ed. 5, p. 647.

BERMUDAS.—Introduced. *Rein.*

Eastern North America, growing in sandy soil, and commoner southward.

[*Panicum maximum*, Jacq. "Guinea grass" has been introduced.]

**Setaria glauca**, Beauv.

*Setaria glauca*, Beauv.; Griseb., Fl. Brit. W. Ind., p. 554; Chapm., Fl. Southern U.S., p. 578.

BERMUDAS.—Introduced. *Moseley.*

In nearly all temperate and tropical countries.

**Oplismenus setarius**, Rœm. et Schult.

*Oplismenus setarius*, Rœm. et Schult., Syst. Veg., ii. p. 481; Kunth, Enum., i. 1, p. 139.

*Orthopogon setarius*, Spreng.; Griseb., Fl. Brit. W. Ind., p. 545.

BERMUDAS.—Introduced. The caves—*Lefroy.*

West Indies, Tropical America, and tropics of the Old World.

**Cenchrus tribuloides**, Linn.

*Cenchrus tribuloides*, Linn., Sp. Pl., ed. 1, p. 1050; Griseb., Fl. Brit. W. Ind., p. 556; Chapm., Fl. Southern U.S., p. 579.

BERMUDAS.—Indigenous. Common on sand hills by the sea—*Lefroy*; the principal binding grass on the sand dunes—*Moseley.*