

**Sporobolus virginicus, Kunth.**

*Sporobolus virginicus*, Kunth, Enum., i. p. 210; Griseb., Fl. Brit. W. Ind., p. 533; Chapm., Fl. Southern U.S., p. 550.

BERMUDAS.—Indigenous. Binding grass on the sand dunes—*Moseley*.

Almost as widely distributed as *Sporobolus indicus*.

*Sporobolus pungens*, Kunth (*errore purgans*), of Rein's list, is probably one of the foregoing species. We have seen no American specimens of the true *Sporobolus pungens*.

**Cynodon dactylon, Pers.**

*Cynodon dactylon*, Pers., Ench., i. p. 85; Griseb., Fl. Brit. W. Ind., p. 540; Chapm., Fl. Southern U.S., p. 557; Kunth, Enum., i. 1, p. 259.

*Agrostis bermudiana*, Tussac ex Kunth, Enum., i. p. 259.

BERMUDAS.—Introduced. Common in cultivated ground—*Lefroy*; *Munro*.

In tropical and temperate countries all over the globe, but probably not indigenous in America.

This is the plant called "Bermuda Grass" in the United States and elsewhere.

**Chloris petræa, Thunb.**

*Chloris petræa*, Thunb., Fl. Cap., p. 109; Griseb., Fl. Brit. W. Ind., p. 539.

*Eustachys petræa*, Desv., Jour. Bot., 1813, i. p. 69; Chapm., Fl. Southern U.S., p. 557.

BERMUDAS.—Indigenous. Common in dry places—*Moseley*; *Jones*; *Reade*.

A coast plant, ranging from North Carolina to Uruguay, and also occurring in Southern Africa.

**Leptochloa mucronata, Kunth.**

*Leptochloa mucronata*, Kunth, Enum., i. 1, p. 270; Griseb., Fl. Brit. W. Ind., p. 537 (excl. synon.); Chapm., Fl. Southern U.S., p. 558.

*Eleusine mucronata*, Michx., Fl. Bor.-Am., i. p. 65.

BERMUDAS.—Introduced. *Rein.*

Virginia and Illinois to Florida and the West Indies; common in cultivated ground.

**Kœleria phleoides, Pers.**

*Kœleria phleoides*, Pers., Ench., i. p. 97; Kunth, Enum., i. p. 383.

BERMUDAS.—Introduced. *Moseley*.

Widely spread in the Old World; but not, as far as we know, naturalised in any part of America.