

Radula pallens, Nees.*Radula pallens*, Nees in G. L. et N. Synop. Hepat., pp. 256 et 726.*Jungermannia pallens*, Sw., Prodr. Fl. Ind. Occ., p. 143, et Fl. Ind. Occ., iii. p. 1847.

BERMUDAS.—In large spreading patches of the usual brown colour, but without perianths. Specimens were also collected in the same state by General Lefroy.

Aneura palmata, Nees ?*Jungermannia palmata*, Hedw. ? Theor., Gen., ed. 1, t. 18, ex G. L. et N. Synop. Hepat., p. 499 (sub *Aneura*).

BERMUDAS.—Without fructification, and in a poor state.

Ottonia aitonia, Lindb. et Nees (*Plagiochasma*).*Plagiochasma aitonia*, Lindb. et Nees in G. L. et N. Synop. Hepat., p. 520.

BERMUDAS.—Barren.

Dumortiera hirsuta, R. Bl. et Nees.*Dumortiera hirsuta*, R. Bl. et Nees in G. L. et N. Synop. Hepat., p. 543.*Marchantia hirsuta*, Sw., Prodr. Fl. Ind. Occ., p. 145, et Fl. Ind. Occ., iii. p. 1879.

BERMUDAS.—Specimens with perfected fruit.

LICHENES.¹
COLLEMACEI.**COLLEMEI.****Leptogium diaphanum**, Mont.*Leptogium diaphanum*, Mont. ; Nyl., Synop. Lich., i. p. 125 ; Crombie in Journ. Linn. Soc. Lond., xvi. p. 214.*Leptogium tremelloides*, var. *mesotomum*, Stirton in Journ. Linn. Soc. Lond., xiv. p. 369.*Lichen diaphanus*, Swartz, Fl. Ind. Occ., p. 1893.

BERMUDAS.—On the bark of trees ; fertile—*Moseley*.

Jamaica ; Peru ; Tahiti ; Philippines.

Leptogium tremelloides, Fries.*Leptogium tremelloides*, Fries ; Nyl., Synop. Lich., i. p. 124 ; Crombie in Journ. Linn. Soc. Lond., xvi. p. 214.*Leptogium marginellum*, Stirton in Journ. Linn. Soc. Lond., xiv. p. 369.

BERMUDAS.—On the bark of trees ; fertile—*Moseley*.

Almost cosmopolitan, except in the colder regions.

¹ The Rev. J. M. Crombie kindly gave his help in drawing up this list.