

LICHENACEI.

CLADONIEI.

Cladonia acuminata, var. *hebescens*, Nyl.

Cladonia acuminata, var. *hebescens*, Nyl., n. var. in Journ. Linn. Soc. Lond., xvi. p. 214.

Differt a *Cladonia acuminata* squamulis crassiusculis subconvexulis. Sterilis modo visa. Forsan propria species.

BERMUDAS.—On the ground ; very sparingly seen—*Moseley*.

Cladonia fibula, Hoffm.

Cladonia fibula, Hoffm., Fl. Germ., p. 127 ; Crombie in Journ. Linn. Soc. Lond., xvi. p. 214.

Cladonia turgida et var. *elegans*, Stirton in Journ. Linn. Soc. Lond., xiv. p. 370.

BERMUDAS.—On the ground ; fertile—*Moseley*.

Europe, and doubtless other regions.

Cladonia mitrula, Tuckerm.

Cladonia mitrula, Tuckerm. in Darl. Fl. Cestr., p. 444, et Gen. Lich., p. 147 ; Nyl., Synop. Lich., i. p. 203 ; Crombie in Journ. Linn. Soc. Lond., xvi. p. 214.

Cladonia botrytes, Stirton in Journ. Linn. Soc. Lond., xiv. p. 370.

BERMUDAS.—On the ground ; fertile—*Moseley*.

Common throughout the southern states of North America, and extending to Mexico and Cuba.

Cladonia pungens, Flöck.

Cladonia pungens, Flöck., Comm. Clad., p. 156 ; Crombie in Journ. Linn. Soc. Lond., xx. p. 83.

Cladonia furcata, Hoffm., var. *pungens*, Fries, Lich. Europ., p. 230 ; Nyl., Synop. Lich., i. p. 207.

BERMUDAS.—On the ground ; sterile—*Moseley*.

Europe ; Africa ; North and South America ; India ; Australia.

RAMALINEI.

Ramalina complanata, Achar.

Ramalina complanata, Achar., Lich. Univ., p. 599 ; Crombie in Journ. Linn. Soc. Lond., xvi. p. 215.

Ramalina calicaris, forma 5, *complanata*, Nyl., Synop. Lich., i. p. 295.

Lichen complanatus, Swartz, Fl. Ind. Occ., p. 1911.

Ramalina bermudiana, Stirton in Journ. Linn. Soc. Lond., xiv. p. 370.

BERMUDAS.—On branches of trees ; fertile—*Moseley*.

West Indies ; Mexico ; subtropical South America ; Africa (Senegal) ; South Australia.