

Lecanora chlaronella, Nyl.*Lecanora chlaronella*, Nyl., n. sp.*Lecanora varia*, var. *symmicta*, Stirton in Journ. Linn. Soc. Lond., xiv. p. 370.

Similis *Lecanoræ chlaroterodi*, Nyl. in Flora, 1876, p. 508, Cubanæ, sed differt gelatina hymenialis iodo vinose fulvescente (præcedentia cærulescentia). Spermogonia non visa.—Journ. Linn. Soc. Lond., xvi. p. 215.

BERMUDAS.—On bark of trees ; a very small specimen only seen—*Moseley*.

Lecanora bermudensis, Nyl.*Lecanora bermudensis*, Nyl., sp. n.*Lecanora subfusca*, var. *argentata*, Stirton in Journ. Linn. Soc. Lond., xiv. p. 371.

Affinis *Lecanoræ umbrinæ*, sed mox differens, margine thallino apotheciorum integro, paraphysibus gracilissentibus, &c. Thallus K.—Journ. Linn. Soc. Lond., xvi. p. 215.

BERMUDAS.—On bark of trees—*Moseley*.

Lecanora glaucmodes, Nyl.*Lecanora glaucmodes*, Nyl. in Flora, 1876, p. 509 ; subsp. *conflectens*, Nyl. in Journ. Linn. Soc. Lond., xvi. p. 215.*Lecanora angulosa*, var. *albo-pruinosa*, Stirton in Journ. Linn. Soc. Lond., xiv. p. 371.

Differt, *Lecanora conflectens* a *Lecanora glaucmodi*, paraphysibus distinctis, iodo gelatina hymeniale vinose fulvescente, &c.

BERMUDAS.—On the bark of trees—*Moseley*.

The typical form was collected in Cuba.

PERTUSARIEI.

Pertusaria papillata, Tuckerm.*Pertusaria papillata*, Tuckerm., Synop. Lich. N. Am., p. 85 ; Journ. Linn. Soc. Lond., xvi. p. 215.*Pertusaria leioplaca*, Stirton in Journ. Linn. Soc. Lond., xiv. p. 371.

BERMUDAS.—On the bark of trees ; a mere fragment—*Moseley*.

North America.

Pertusaria pustulata, Schær.*Pertusaria pustulata*, Schær, Spicil., p. 66 ; Nyl. in Triana et Planchon Prod. Flor. Nov. Granat. Crypt., p. 35 ; Crombie in Journ. Linn. Soc. Lond., xvi. p. 216.*Pertusaria melaleuca*, Turn et Borr. ; Stirton in Journ. Linn. Soc. Lond., xiv. p. 371.

BERMUDAS.—On the bark of trees ; sparingly seen—*Moseley*.

Europe and North and South America.