

Ustilago carbo, Tul.

Ustilago carbo, Tul, Mém. Ustil., p. 78; Fisch. Waldh. in Ann. Sc. Nat., série 6, iv. p. 200; Berkl. in Journ. Linn. Soc. Lond., xv. p. 49.

BERMUDAS.—Without any memorandum—*Moseley*.

Generally dispersed where wheat, oats, and barley are cultivated.

Puccinia lantanæ, Farlow.

Puccinia lantanæ, Farlow, MSS. in Herb. Cooke.

Hypophylla. Uredo cum teleutosporis immixta.

Soris minutis sparsis vel plerumque in cæspitulis maculæformibus aggregatis, atro-fuscis. Uredosporis globosis ovatisque lævibus fuscis (0·02 × 0·015 mm. diam.) Teleutosporis ellipticis, medio constrictis, supra rotundatis, infra in stipitem brevem attenuatis, lævibus, fuscis (0·03–0·035 × 0·018–0·02 mm).—M. C. Cooke, MSS.

BERMUDAS.—On leaves of *Lantana odorata*—*Farlow*.

ASCOMYCETES.

PYRENOMYCETES.

Hypoxylon concentricum, Grev.

Hypoxylon concentricum, Grev., Crypt. Scot. Flor., vi. t. 324; Saccardo, Syll. Fung., iii. p. 393 (sub *Daldinia*); Berkl. in Journ. Linn. Soc. Lond., xv. p. 49.

BERMUDAS.—On dead wood, Devonshire Marsh—*Moseley*.

Widely diffused in temperate and tropical regions.

Hypoxylon multiforme, Fries.

Hypoxylon multiforme, Fries, Summ. Veg. Scand., p. 384; Saccardo, Syll. Fung., i. p. 363; Berkl. in Journ. Linn. Soc. Lond., xiv. p. 351.

BERMUDAS.—On decayed wood—*Moseley*.

Throughout Europe, India, and North America, northward to Kamtschatka.

Sphæria mammæformis, Pers.

Sphæria mammæformis, Pers., Synop. Fung., p. 64; Saccardo, Syll. Fung., iii. p. 258 (sub *Rosellinia*); Berkl. in Journ. Linn. Soc. Lond., xiv. p. 351.

BERMUDAS.—On dead sticks, both on the bark and on the wood; in the former case looking very like a *Verrucaria*—*Moseley*.

North and south temperate regions.