

Taonia schröderi, Mert.

Taonia schröderi, Mert. ex J. G. Agardh, Sp. Alg., i. p. 102; Harv., Nereis Bor.-Am., i. p. 107.

BERMUDAS. *Farlow*.

West Indies; Brazil; India; Ceylon.

Dictyota ciliata, J. G. Agardh.

Dictyota ciliata, J. G. Agardh in Linnæa, 1841, p. 5; Kütz., Sp. Alg., p. 556, non Lamour. in Journ. de Bot. Paris, 1809, ii. p. 41; Harv., Nereis Bor.-Am., i. p. 110; Rein in Bericht Senckenb. Naturf. Gesellsch., Frankf. am M., 1873, p. 151.

BERMUDAS. *Kemp*; *Rein*.

Florida; West Indies; Brazil; Australia; Ceylon.

Dictyota bartayresiana, Lamour.

Dictyota bartayresiana, Lamour. in Journ. de Bot. Paris, 1809, ii. p. 43; Kütz., Sp. Alg., p. 554; Harv., Nereis Bor.-Am., i. p. 110; Dickie in Journ. Linn. Soc. Lond., xiv. p. 313.

BERMUDAS.—Shallows only—*Moseley*

“Small and rather imperfect, but, I think, may be referred to this species.”—*Dickie*.

Florida, West Indies, shores of the Atlantic Islands, the Persian Gulf, and Mauritius.

Dictyota ligulata, Kütz.

Dictyota ligulata, Kütz. in Bot. Zeit., 1847, p. 53, et Sp. Alg., p. 554; Dickie in Journ. Linn. Soc. Lond., xiv. p. 313.

BERMUDAS.—From thirty-one fathoms—*Moseley*.

Mediterranean.

Dictyota dichotoma, Lamour.

Dictyota dichotoma, Lamour. in Journ. de Bot. Paris, 1809, ii. p. 42; Kütz., Sp. Alg., p. 554; Harv., Nereis Bor.-Am., i. p. 109; Dickie in Journ. Linn. Soc. Lond., xiv. p. 313.

Ulva dichotoma, Huds., Fl. Angl., ed. 1, p. 476.

BERMUDAS.—Some from thirty-one fathoms seem to be dwarf forms of this species—*Moseley*; *Kemp*; *Rein*.

Very widely diffused.

Dictyota crenulata, J. G. Agardh.

Dictyota crenulata, J. G. Agardh, Fucoid., p. 94; Kütz., Sp. Alg., p. 558; Harv., Nereis Bor.-Am., i. p. 111; Kemp. in Canad. Nat. and Geol., May 1857.

BERMUDAS. *Kemp*.

Eastern coast of Mexico.