

**Dictyota fasciola, Lamour.**

*Dictyota fasciola*, Lamour. in Journ. de Bot. Paris, 1809, ii. p. 43; Kütz., Sp. Alg., p. 555; Harv., Nereis Bor.-Am., i. p. 108; Rein in Bericht Senckenb. Naturf. Gesellsch., Frankf. am M., 1873, p. 151.

*Zonaria fasciola*, K. A. Agardh, Sp. Alg., i. p. 136.

BERMUDAS. *Rein*.

Florida; Mediterranean; Red Sea; Ceylon.

**Asperococcus sinuosus, Bory.**

*Asperococcus sinuosus*, Bory, Morée, iii. p. 326, ex J. G. Agardh, Sp. Alg., i. p. 75; Harv., Nereis Bor.-Am., i. p. 117; Dickie in Journ. Linn. Soc. Lond., xiv. p. 313.

*Enceclium sinuosum*, K. A. Agardh, Sp. Alg., p. 146; Kütz., Sp. Alg., p. 552.

BERMUDAS.—In shallow water—*Moseley*; *Kemp*; *Rein*.

Widely diffused, though we have seen no specimens from Polynesia or Western America, except from Juan Fernandez and the extreme northwest.

## CHORDARIACEÆ.

**Mesogloia vermicularis, K. A. Agardh.**

*Mesogloia vermicularis*, K. A. Agardh, Syst. Alg., p. 51; Kütz., Sp. Alg., p. 544; Harv., Nereis Bor.-Am. i. p. 126; Rein in Bericht Senckenb. Naturf. Gesellsch., Frankf. am M., 1873, p. 152.

BERMUDAS. *Kemp*; *Rein*.

Western Europe; Mediterranean.

**Mesogloia griffithsiana, Grev.**

*Mesogloia griffithsiana*, Grev.; Kütz., Sp. Alg., p. 545; Harv., Nereis Bor.-Am., i. p. 127; Kemp in Canad. Nat. and Geol., May 1857.

BERMUDAS. *Kemp*.

Coast of England; Heligoland.

Perhaps an error in determination, or the specimen from some other locality.

**Mesogloia virescens, Carm.**

*Mesogloia virescens*, Carm.; Hook., Br. Fl., ii. p. 387, ex Kütz., Sp. Alg., p. 545; Harv., Nereis Bor.-Am., i. p. 126; Dickie in Journ. Linn. Soc. Lond., xiv. p. 313.

BERMUDAS.—Shallows only—*Moseley*; *Kemp*.

Western Europe; eastern North America; Australasia.