

Gelidium corneum, Lamour.

Gelidium corneum, Lamour., Ess., p. 41; Kütz., Sp. Alg., p. 764; J. G. Agardh, Sp. Alg., ii. p. 469; Harv., Phyc. Brit., t. 53; Harv., Nereis Bor.-Am., ii. p. 116; Kemp in Canad. Nat. and Geol., May 1857.

BERMUDAS. *Kemp*.

Generally diffused.

Wurdemannia setacea, Harv.

Wurdemannia setacea, Harv., Nereis Bor.-Am. ii., p. 246; Kütz., Tab. Phyc., xix. t. 26; Rein in Bericht Senckenb. Naturf. Gesellsch., Frankf. am M., 1873, p. 152.

BERMUDAS. *Rein*.

Florida Keys.

Hypnea musciformis, Lamour.

Hypnea musciformis, Lamour., Ess., p. 43; J. G. Agardh, Sp. Alg., ii. p. 442; Kütz., Sp. Alg., p. 758; Harv., Nereis Bor.-Am., ii. p. 123; Rein in Bericht Senckenb. Naturf. Gesellsch., Frankf. am M., 1873, p. 152.

BERMUDAS. *Kemp*; *Rein*.

Mediterranean; Atlantic; Indian Ocean; Australasia, &c.

Hypnea cornuta, Lamour.

Hypnea cornuta, Lamour., ex J. G. Agardh, Sp. Alg., ii. p. 449; Harv., Nereis Bor.-Am., ii. p. 125; Dickie in Journ. Linn. Soc. Lond., xiv. p. 314.

BERMUDAS.—From shallows and from thirty-one fathoms—*Moseley*.

Florida; West Indies; Guiana; Manilla; Ceylon; Mauritius.

Eucheuma isiforme, J. G. Agardh.

Eucheuma isiforme, J. G. Agardh, Sp. Alg., ii. p. 627; Harv., Nereis Bor.-Am., ii. p. 118; Dickie in Journ. Linn. Soc. Lond., xiv. p. 314.

BERMUDAS. *Moseley*; *Kemp*; *Rein*.

Florida; West Indies.

SQUAMARIEÆ.

Peyssonelia dubyi, Crouan.

Peyssonelia dubyi, Crouan in Ann. Sc. Nat., 1844, p. 368, t. 11, B.; Harv., Phyc. Brit., t. 71, et Nereis Bor.-Am., ii. p. 130; Kütz., Sp. Alg., p. 694; Dickie in Journ. Linn. Soc. Lond., xiv. p. 314.

BERMUDAS.—From thirty-one fathoms—*Moseley*.

Coast of Britain and Northern France; Cape Verde Islands; Florida.