

HELMINTHOCLADEÆ.

*Liagora valida*, Harv.

*Liagora valida*, Harv., *Nereis Bor.-Am.*, ii. p. 138, t. 31, A.; Dickie in *Journ. Linn. Soc. Lond.*, xiv. p. 314.

BERMUDAS. In shallows—*Moseley*; *Kemp*; *Rein.*

Florida Keys.

The plant named *Liagora pulverulenta* by Kemp may be the same as this.

*Galaxaura fastigiata*, Dcne.

*Galaxaura fastigiata*, Dcne.; Kütz., Sp. Alg., p. 530; Rein in *Bericht Senckenb. Naturf. Gesellsch.*, Frankf. am M., 1873, p. 152.

BERMUDAS. *Rein.*

Philippines.

We include this entirely on Rein's authority, having seen only Cuming's specimens from the Philippines.

*Galaxaura lapidescens*, Lamour.

*Galaxaura lapidescens*, Lamour., *Polyp. Flex.*, p. 264; Kütz., Sp. Alg., p. 530; Dickie in *Journ. Linn. Soc. Lond.*, xiv. p. 315.

BERMUDAS.—From thirty fathoms—*Moseley*.

Cape Verde and Canary Islands; St Helena; Florida; Sandwich Islands; Polynesia; Mauritius.

*Galaxaura rugosa*, Lamour.

*Galaxaura rugosa*, Lamour., *Polyp. Flex.*, p. 263; Kütz., Sp. Alg., p. 530; Dickie in *Journ. Linn. Soc. Lond.*, xiv. p. 315.

BERMUDAS.—From shallows, and also from thirty-one fathoms—*Moseley*.

Cape Verde Islands; West Indies; Mauritius.

*Scinaia furcellata*, Bivona.

*Scinaia furcellata*, Bivona in L'Iride, Palermo, 1822, cum icono; J. G. Agardh, Sp. Alg., ii. p. 422; Harv., *Nereis Bor.-Am.*, ii. p. 136; Dickie in *Journ. Linn. Soc. Lond.*, xiv. p. 315.

BERMUDAS.—A solitary and imperfect specimen from thirty-one fathoms—*Moseley*. Widely diffused.