

**Helminthora divaricata**, J. G. Agardh.

*Helminthora divaricata*, J. G. Agardh, Sp. Alg., ii. p. 416; Harv., Nereis Bor.-Am., ii. p. 133; Kemp in Canad. Nat. and Geol., May 1857.

BERMUDAS. *Kemp*.

Florida; Western Europe; Mediterranean; Australasia.

## RHODYMENIACEÆ.

**Rhodymenia palmata**, Grev.

*Rhodymenia palmata*, Grev.; Harv., Phyc. Brit., tt. 217 et 218, et Nereis Bor.-Am., ii. p. 148; Kemp in Canad. Nat. and Geol., May 1857.

BERMUDAS. *Kemp*.

A common and widely dispersed species; yet neither this nor the next was collected in the Bermudas by either Rein or Moseley.

## SPYRIDIACEÆ.

**Spyridia aculeata**, Kütz.

*Spyridia aculeata*, Kütz., Sp. Alg., p. 668 (excl. synonym.), ex Harv., Nereis Bor.-Am., ii. p. 205; Kemp in Canad. Nat. and Geol., May 1857.

BERMUDAS. *Kemp*.

Florida Keys; Cadiz; Cape Verde Islands; Mediterranean; Red Sea.

**Spyridia filamentosa**, Harv.

*Spyridia filamentosa*, Harv., Nereis Bor.-Am., ii. p. 204, t. 34, A.; Rein in Bericht Senckenb. Naturf. Gesellsch., Frankf. am M., 1873, p. 152.

BERMUDAS. *Rein*.

Eastern North America, from Massachusetts to Florida.

## CRYPTONEMIACEÆ.

**Callophyllis laciniata**, Kütz.

*Callophyllis laciniata*, Kütz., Sp. Alg., p. 744; Harv., Nereis Bor.-Am., ii. p. 171.

*Rhodymenia laciniata*, Grev.; Harv., Phyc. Brit., t. 121; Kemp in Canad. Nat. and Geol., May 1857.

BERMUDAS. *Kemp*.

Western Europe; Mediterranean; California; Cape Heulopen, eastern North America.

**Kallymenia reniformis**, J. G. Agardh.

*Kallymenia reniformis*, J. G. Agardh., Sp. Alg., ii. p. 286? Dickie in Journ. Linn. Soc. Lond., xiv. p. 315.

BERMUDAS.—A few very fragmentary specimens from thirty-one fathoms—*Moseley*.

The typical plant inhabits the shores of Western Europe.