

Chrysomenia halymenioides, Harv.

Chrysomenia halymenioides, Harv., Nereis Bor.-Am., ii. p. 188, t. 20, A.; Dickie in Journ. Linn. Soc. Lond., xiv. p. 315.

BERMUDAS.—A single very small example one inch high—*Moseley*.

Florida Keys.

Harvey states that this is very closely allied to the next.

Chrysomenia dichotoma, J. G. Agardh?

Chrysomenia dichotoma, J. G. Agardh, Sp. Alg., ii. p. 211? Dickie in Journ. Linn. Soc. Lond., xiv. p. 315; Harv., Nereis Bor.-Am., ii. p. 189.

BERMUDAS.—From thirty-one fathoms—*Moseley*.

Adriatic; Mediterranean; Teneriffe.

Chylocladia rosea, Harv.

Chylocladia rosea, Harv., Phyc. Brit., t. 301, et Nereis Bor.-Am., p. 186; Kemp in Canad. Nat. and Geol., May 1857.

BERMUDAS. *Kemp*.

Eastern coast of North America and Western Europe.

We have seen no specimens from the American side of the Atlantic farther south than Rhode Island.

Catenella pinnata, Harv.

Catenella pinnata, Harv., Nereis Bor.-Am., ii. p. 201, t. 29, B.

BERMUDAS. *Farlow*.

Florida Keys.

Gloiosiphonia capillaris, Carm.

Gloiosiphonia capillaris, Carm.; J. G. Agardh, Sp. Alg., ii. p. 161; Harv., Nereis Bor.-Am. ii. p. 202; Kemp in Canad. Nat. and Geol., May 1857.

BERMUDAS. *Kemp*.

Western Europe from Norway to Spain; eastern North America, on the coast of New England; Vancouver Island, Pacific.

CERAMIACEÆ.

Centroceras clavulatum, J. G. Agardh.

Centroceras clavulatum, J. G. Agardh, Sp. Alg., ii. p. 148; Harv., Nereis Bor.-Am., ii. p. 211, t. 33, C.; Dickie in Journ. Linn. Soc. Lond., xiv. p. 315.

BERMUDAS.—Shallow water—*Moseley*; *Rein*.