

**Caulerpa plumaris**, K. A. Agardh.

*Caulerpa plumaris*, K. A. Agardh, Sp. Alg., i. p. 436; Kütz., Sp. Alg., p. 496; Harv., Nereis Bor.-Am., iii. p. 17, t. 38, C.; Rein in Bericht Senckenb. Naturf. Gesellsch., Frankf. am M., 1873, p. 153.

BERMUDAS. *Kemp*; *Rein*; *Farlow*.

Florida Keys, and generally dispersed in warm seas.

**Caulerpa clavifera**, K. A. Agardh.

*Caulerpa clavifera*, K. A. Agardh, Sp. Alg., i. p. 437; Harv., Nereis Bor.-Am., iii. p. 19; Dickie in Journ. Linn. Soc. Lond., xiv. p. 315.

*Chanoinia clavifera*, Kütz., Sp. Alg., p. 498.

BERMUDAS.—In lagoons—*Moseley*; *Rein*.

Also a very widely diffused species.

**Caulerpa prolifera**, Lamour.

*Caulerpa prolifera*, Lamour. in Journ. de Bot., Paris, 1809, ii. p. 136; Harv., Nereis Bor.-Am., iii. p. 16, t. 38, B.; Rein in Bericht Senckenb. Naturf. Gesellsch., Frankf. am M., 1873, p. 153.

*Phyllerpa prolifera*, Kütz., Sp. Alg., p. 494.

BERMUDAS. *Kemp*; *Rein*.

Widely distributed in the North Atlantic and Mediterranean.

**Caulerpa taxifolia**, K. A. Agardh.

*Caulerpa taxifolia*, K. A. Agardh, Sp. Alg., i. p. 435, cum  $\beta$  *crassifolia*, l. c., p. 436; Kütz., Sp. Alg., p. 495.

BERMUDAS.—Walsingham—*Farlow*.

Widely dispersed in warm seas.

**Halimeda incrassata**, Lamour.

*Halimeda incrassata*, Lamour., Exp. Méth., p. 26, ex Kütz., Sp. Alg., p. 504; Harv., Nereis Bor.-Am., iii. p. 24; Dickie in Journ. Linn. Soc. Lond., xv. p. 488; Rein in Bericht Senckenb. Naturf. Gesellsch., Frankf. am M., 1873, p. 153.

BERMUDAS. *Moseley*; *Rein*.

West Indies; Florida Keys; North Australia; Friendly Islands.

**Halimeda opuntia**, Lamour.

*Halimeda opuntia*, Lamour., Exp. Méth., p. 27, t. 20, fig. 6; Kütz., Sp. Alg., p. 504; Harv., Nereis Bor.-Am., iii. p. 23, t. 40, B.; Rein in Bericht Senckenb. Naturf. Gesellsch., Frankf. am M., 1873, p. 153.

BERMUDAS. *Rein*.