

Cladophora luteola, Harv.

Cladophora luteola, Harv., Nereis Bor.-Am., iii. p. 81; Dickie in Journ. Linn. Soc. Lond., xiv. p. 316.

BERMUDAS. *Moseley*; *Rein*.

The identity of Moseley's specimens with this species is somewhat doubtful.

Cladophora lutescens, Kütz.

Cladophora lutescens, Kütz., Phyc. Germ., p. 211, et Sp. Alg., p. 403; Rein in Bericht Senckenb. Naturf. Gesellsch., Frankf. am M., 1873, p. 153.

BERMUDAS. *Rein*.

North Sea; eastern North Atlantic; Mediterranean.

Cladophora prolifera, Kütz.

Cladophora prolifera, Kütz., Phyc. Germ., p. 207, et Sp. Alg., p. 390; Dickie in Journ. Linn. Soc. Lond., xiv. p. 316.

BERMUDAS. *Moseley*.

Mediterranean; eastern North Atlantic; Brazil; Peru; Bombay; Polynesia.

Hormotrichum bermudianum, Harv.

Hormotrichum bermudianum, Harv., sp. n., ex Rein in Bericht Senckenb. Naturf. Gesellsch., Frankf. am M., 1873, p. 153 (*absque descriptione*).

BERMUDAS. *Rein*.

We have seen no specimen of this species.

Chætomorpha ærea, Kütz.

Chætomorpha ærea, Kütz., Sp. Alg., p. 379; Harv., Nereis Bor.-Am., iii. p. 86; Dickie in Journ. Linn. Soc. Lond., xiv. p. 316.

Conferva ærea, Dillw., t. 80; Harv., Phyc. Brit., t. 99.

BERMUDAS.—In shallow water—*Moseley*.

Widely spread.

Chætomorpha geniculata, Mont.

Chætomorpha geniculata, Mont., Syll. Gen. et Sp. Crypt., p. 460; Rein in Bericht Senckenb. Naturf. Gesellsch., Frankf. am M., 1873, p. 153.

BERMUDAS. *Rein*.

West Indies.