

COMBRETACEÆ.

Laguncularia racemosa, Gært.

Laguncularia racemosa, Gært. ; Griseb., Fl. Brit. W. Ind., p. 276 ; Chapman, Fl. Southern U.S., p. 136 (*Conocarpus racemosus*, Linn.)

On the shore. A maritime shrub from Florida to Bahia, and on the western coast of Africa.

MYRTACEÆ.

Eugenia axillaris, Poir., is doubtless the same that we have referred to *Eugenia monticola*, DC.

RUBIACEÆ.

Spermacoce tenuior, Lam. In the Walsingham tract.

BORAGINEÆ.

Lithospermum distichum, Orteg.

Lithospermum distichum, Orteg. ; DC., Prodr., x. p. 77.

This Mexican plant is enumerated by Sir J. H. Lefroy as a native of the sea-shore.

VERBENACEÆ.

Callicarpa ferruginea, Swartz ?

Callicarpa ferruginea, Swartz ? Griseb., Fl. Brit. W. Ind., p. 499.

Chiefly in the Walsingham tract. A common West Indian species.

URTICACEÆ.

Sponia lamarckiana, Dcne.

Sponia lamarckiana, Dcne. ; Griseb., Fl. Brit. W. Ind., p. 150.

Forms the underwood of a large part of the Walsingham tract, especially near Paynter's Vale. A common West Indian species.

JUNCACEÆ.

Juncus maritimus, Lam.

Juncus maritimus, Lam. ; Chapman, Fl. Southern U.S., p. 493.

Common in the wetter portions of the marshes. A widely dispersed rush.

GRAMINEÆ.

Arundinaria tecta, Muhl.

Arundinaria tecta, Muhl. ; Chapman, Fl. Southern U.S., p. 561.

This common swamp reed of the south-eastern States is recorded as indigenous in the Bermudas, but no special locality is given.