

ENUMERATION OF THE PLANTS.

PHANEROGAMÆ.—DICOTYLEDONES.

CAPPARIDEÆ.

Capparis cynophallophora, Linn.

Capparis cynophallophora, Linn., Sp. Pl., ed. 1, p. 504; DC., Prodr., i. p. 249; Jacq., Sel. Stirp. Amer., t. 98; Griseb., Fl. Brit. W. Ind., p. 18 (varieties).

FERNANDO-NORONHA. *Middleton*; *Darwin*; *Moseley*.

Very widely dispersed in Tropical America, especially in maritime districts.

PORTULACEÆ.

Portulaca oleracea, Linn.

Portulaca oleracea, Linn., Sp. Pl., ed. 1, p. 445; DC., Prodr., iii. p. 353; Griseb., Fl. Brit. W. Ind., p. 57.

FERNANDO-NORONHA. *Moseley*.

In nearly all tropical and subtropical countries, especially in maritime districts.

Talinum patens, Willd.

Talinum patens, Willd., Sp. Pl., ii. p. 863; Jacq., Hort. Vindob., ii. t. 151.

Portulaca paniculata, Griseb., Fl. Brit. W. Ind., p. 57.

FERNANDO-NORONHA. *Moseley*.

In maritime districts chiefly, from Mexico and the West Indies to Buenos Ayres.

MALVACEÆ.

Sida spinosa, Linn.

Sida spinosa, Linn., Sp. Pl., ed. 1, p. 683; Griseb., Fl. Brit. W. Ind., p. 74.

FERNANDO-NORONHA. *Moseley*.

Very common in the warmer parts of America, and generally dispersed in the tropics of the Old World.