

SAPINDACEÆ.

Cardiospermum halicacabum, Linn.

Cardiospermum halicacabum, Linn., Sp. Pl., ed. 1, p. 366 ; Griseb., Fl. Brit. W. Ind., p. 122.

FERNANDO-NORONHA. *Moseley*.

Common in tropical and subtropical countries all over the globe.

ANACARDIACEÆ.

Anacardium occidentale, Linn.

Anacardium occidentale, Linn., Sp. Pl., ed. 1, p. 383 ; Griseb., Fl. Brit. W. Ind., p. 176 ; Desc., Fl. Ant., vii. t. 507.

FERNANDO-NORONHA. *Moseley*.

Mexico and West Indies to Brazil, and naturalised in nearly all other tropical countries.

LEGUMINOSÆ.

PAPILIONACEÆ.

Tephrosia cinerea, Pers.

Tephrosia cinerea, Pers., Ench., ii. p. 328 ; Jacq., Ic. Pl. Par., t. 575 ; Griseb., Fl. Brit. W. Ind. p. 182.

FERNANDO-NORONHA. *Moseley*.

Mexico and the West Indies to Uruguay.

Phaseolus peduncularis, H. B. K.

Phaseolus peduncularis, H. B. K., Nov. Gen. et. Sp., vi. p. 447 ; Benth. in Mart. Fl. Bras., xv. pars 1, p. 184.

FERNANDO-NORONHA. *Moseley*.

Panama eastward, and southward to North Brazil.

Desmodium spirale, DC.

Desmodium spirale, DC., Prodr., ii. p. 332 ; Benth. in Mart. Fl. Bras., xv. pars 1, p. 105 ; Griseb., Fl. Brit. W. Ind., p. 188.

FERNANDO-NORONHA. *Moseley*.

Common in tropical regions in both hemispheres.

CÆSALPINIÆ.

Cassia tora, Linn.

Cassia tora, Linn., Sp. Pl., ed. 1, p. 376 ; Benth. in Trans. Linn. Soc. Lond., xxvii. p. 535 ; Griseb., Fl. Brit. W. Ind., p. 209, sub *Cassia obtusifolia*.

FERNANDO-NORONHA. *Moseley*.

Generally dispersed in the tropics.