

Cucumis anguria, Linn.

Cucumis anguria, Linn., Sp. Pl., ed. i. p. 1011; Cogniaux in DC. Monogr. Phanerog., iii. p. 501;
Griseb., Fl. Brit. W. Ind., p. 288.

ST MICHAEL'S MOUNT. *Moseley*.

Florida and Texas, through Mexico and the West Indies to Peru and Brazil.

Momordica charantia, Linn.

Momordica charantia, Linn., Sp. Pl., ed. i. p. 1009; Cogniaux in DC. Monogr. Phanerog., iii. p. 436;
Griseb., Fl. Brit. W. Ind., p. 287.

RAT ISLAND and ST MICHAEL'S MOUNT. *Moseley*.

Everywhere in tropical and subtropical regions, but supposed to have been introduced into America.

RUBIACEÆ.

Spermacoce parviflora, Hemsl.

Spermacoce parviflora, Hemsl. in Salv. et Godm. Biolog. Centr. Am., Bot., ii. p. 59.
Borreria parviflora, G. F. W. Mey., Fl. Esseq., p. 83, t. i.; Griseb., Fl. Brit. W. Ind., p. 349.
Spermacoce hirta, Swartz non alior.

FERNANDO-NORONHA. *Moseley*.

Common in Tropical America, from Guatemala to Peru and Brazil, and throughout the West Indies.

COMPOSITÆ.

Blainvillea rhomboidea, Cass.

Blainvillea rhomboidea, Cass. in Dict. Sc. Nat., xxix. p. 494; DC., Prodr., v. p. 492; Oliv., Fl. Trop. Afr., iii. p. 375.

FERNANDO-NORONHA. *Moseley*.

Common in Brazil; also in Tropical Africa.

Eclipta alba, Hassk.

Eclipta alba, Hassk., Pl. Jav. Rar., p. 528; Griseb., Fl. Brit. W. Ind., p. 370.
Eclipta erecta, *Eclipta prostrata*, et *Eclipta punctata*, Linn., Mant. alt., p. 286.

FERNANDO-NORONHA. *Moseley*.

A common weed in nearly all tropical and subtropical countries, and extending into temperate regions.