

APOCYNACEÆ.

Rauwolfia ternifolia, Kunth.

Rauwolfia ternifolia, Kunth, Synop. Pl. Æquin., ii. p. 298; Griseb., Fl. Brit. W. Ind., p. 408; Bot. Mag., t. 2440.

FERNANDO-NORONHA. *Moseley*.

Trinidad, Venezuela, Guiana, and Brazil.

ASCLEPIADEÆ.

Gonolobus micranthus, Hook. (Plate XV.)

Gonolobus micranthus, Hook. f. MSS. in Herb. Kew.

Gracilis, floribus pro genere minutis, folliculis grosse muricato-tuberculatis.

Frutex volubilis vel scandens, ramulis gracilibus puberulis, cortice flavo. *Folia* longe graciliterque petiolata, membranacea, cordiformia, sinu clauso, lobis sæpius obtegentibus, absque petiolo 1-2 poll. longa, acuta, utrinque sparse beviderque pubescentia, supra in sinu 2-3 glandulosa; petiolus 1-2 poll. longus. *Flores* minuti, pauci in axillas conferti; calyx intus 5-glandulosus, lobis ovatis acutis intus glabris; corolla rotato-campanulata, intus setuloso-pubescentia, lobis ovatis acutiusculis apicem versus tantum hirsutis; corona annulata, puberula, intus lineis crassis instructa, 5-lobata, lobis corollæ segmentis alternantibus gynostegio brevioribus; stigma planum. *Folliculi* longe grosseque muricato-tuberculati, simul furfuraceo-hirsuti; semina longe comosa, ovata, compressa, margine muricata.

ST MICHAEL'S MOUNT. *Moseley*.

BORAGINEÆ.

Heliotropium indicum, Linn.

Heliotropium indicum, Linn., Sp. Pl., ed. 1, p. 130; Griseb., Fl. Brit. W. Ind., p. 485; Bot. Mag. t. 1837.

Heliophytum indicum, DC., Prodr., ix. p. 556.

FERNANDO-NORONHA. *Darwin*; *Moseley*.

A common weed in warm countries.

CONVOLVULACEÆ.

Ipomœa pentaphylla, Jacq.

Ipomœa pentaphylla, Jacq., Collect., ii. p. 297, et Ic. Pl. Rar., t. 319; Griseb., Fl. Brit. W. Ind., p. 467.

Batatas pentaphylla, Choisy in DC. Prodr., ix. p. 339.

FERNANDO-NORONHA. *Moseley*.

Generally dispersed in tropical countries.