

Ipomoea pes-caprae, Sweet.

Ipomoea pes-caprae, Sweet, Hort. Suburb. Londin. (1818), p. 35; Roth, Nov. Pl. Sp., p. 100; Griseb., Fl. Brit. W. Ind., p. 470.

FERNANDO-NORONHA. *Moseley*.

Very widely dispersed in warm countries, but chiefly on sandy sea-shores.

Ipomoea tuba, Don.

Ipomoea tuba, Don, Gen. Syst., iv. p. 271; Griseb., Fl. Brit. W. Ind., p. 467; Mart., Fl. Bras., vii. p. 216.

FERNANDO-NORONHA. *Moseley*.

West Indies, Guiana, and Brazil.

Cuscuta americana, Linn.

Cuscuta americana, Linn., Sp. Pl., ed 1, p. 124; Griseb., Fl. Brit. W. Ind., p. 476; Mart., Fl. Bras., vii. p. 376, t. 126, fig. 1.

FERNANDO-NORONHA. *Moseley*.

Mexico and the West Indies to Brazil.

SOLANACEÆ.

Solanum paniculatum, Linn.

Solanum paniculatum, Linn., Sp. Pl., ed. 2, p. 267; DC., Prodr., xiii. 1, p. 278.

FERNANDO-NORONHA. *Moseley*.

Common in Brazil.

Physalis hirsuta, Dun. var. ?

Physalis hirsuta, Dun. var. ? DC., Prodr., xiii. 1, p. 445.

FERNANDO-NORONHA. *Moseley*.

This may be the same as *Physalis pubescens*, Linn., a widely dispersed species in warm countries.

Datura stramonium, Linn.

Datura stramonium, Linn., Sp. Pl., ed. 1, p. 179; Griseb., Fl. Brit. W. Ind., p. 434.

FERNANDO-NORONHA. *Moseley*.

Everywhere, almost, in tropical and temperate regions where there is cultivation.

SCROPHULARINEÆ.

Scoparia dulcis, Linn.

Scoparia dulcis, Linn., Sp. Pl., ed. 1, p. 116; Griseb., Fl. Brit. W. Ind., p. 427; DC., Prodr., x. p. 431.

FERNANDO-NORONHA. *Moseley*.

Common in nearly all tropical and subtropical countries.