

HELMINTHOCLADEÆ.

Galaxaura cylindrica, Lamour.

Galaxaura cylindrica, Lamour., Exp. Méth., p. 22, ex Kütz., Sp. Alg., p. 530; Dickie in Journ. Linn. Soc. Lond., xiv. p. 364.

FERNANDO-NORONHA. *Moseley.*

Barbadoes; Red Sea; Cape Verde Islands.

Galaxaura oblongata, Lamour.

Galaxaura oblongata, Lamour., Polyp. Flex., p. 262, ex Kütz., Sp. Alg., p. 529; Dickie in Journ. Linn. Soc. Lond., xiv. p. 364.

FERNANDO-NORONHA. *Moseley.*

West Indies; Ceylon; Cape of Good Hope.

Galaxaura marginata, Lamour.

Galaxaura marginata, Lamour.; Kütz., Sp. Alg., p. 530; Dickie in Journ. Linn. Soc. Lond., xiv. p. 364.

FERNANDO-NORONHA. *Moseley.*

West Indies; Madagascar; Mauritius.

Galaxaura rugosa, Lamour.

Galaxaura rugosa, Lamour.; Kütz., Sp. Alg., p. 530; Dickie in Journ. Linn. Soc. Lond., xiv. p. 364.

FERNANDO-NORONHA. *Moseley.*

West Indies; Mauritius; Gambier Islands; Cape Verde Islands; Fiji Islands.

CRYPTONEMIACEÆ.

Chrysymenia enteromorpha, Harv. ?

Chrysymenia enteromorpha, Harv., Nereis Bor.-Am., ii. p. 187? ex Dickie in Journ. Linn. Soc. Lond., xiv. p. 364.

FERNANDO-NORONHA. *Moseley.*

Florida.

The single very fragmentary specimen has the structure of the genus, but the species is doubtful.

CHLOROSPERMEÆ.

SIPHONACEÆ.

Caulerpa clavifera, K. A. Agardh.

Caulerpa clavifera, K. A. Agardh, Sp. Alg., i. p. 437; Harv., Nereis Bor.-Am., iii. p. 19; Dickie in Journ. Linn. Soc. Lond., xiv. p. 364.

FERNANDO-NORONHA. *Moseley.*

Canaries; Red Sea; Indian Ocean; Florida Keys.