

RAMALINEI.

Ramalina pollinaria, Acharius.

Ramalina pollinaria, Acharius, Lich. Univ., p. 608; A. Rich., Sertum Astrolab., p. 51; Nyl., Synop. Lich., i. p. 296.

ASCENSION. *Lesson.*

Europe; Asia; Africa; West Indies; South America; St Helena.

This is included on the authority of Richard.

Ramalina polymorpha, Acharius.

Ramalina polymorpha, Acharius, Lich. Univ., p. 600, Synop. Lich., p. 295; Nyl., Synop. Lich., i. p. 293.

Ramalina implicata (errore pro *empecta*, Acharius?) A. Rich., Sertum Astrolab., p. 51.

ASCENSION. *Lesson.*

Widely dispersed.

USNEEI.

Usnea trichodea, Acharius.

Usnea trichodea, Acharius, Meth. Lich., p. 312, t. 8, fig. 1; Nyl., Synop. Lich., i. p. 270; A. Rich., Sertum Astrolab., p. 50.

ASCENSION. *Lesson.*

Nova Scotia; Cuba; Cape of Good Hope; Bourbon; New Zealand; Java.

PARMELIEI.

Parmelia conoplea, Acharius.

Parmelia conoplea, Acharius, Synop. Lich., p. 213, ex A. Rich., Sertum Astrolab., p. 51.

ASCENSION. *Lesson.*

Parmelia saxatilis, Acharius.

Parmelia saxatilis, Acharius, Synop. Lich., p. 203; Nyl., Synop. Lich., i. p. 388; A. Rich., Sertum Astrolab., p. 51.

ASCENSION. *Lesson.*

Generally diffused.

PHYSCIEL.

Physcia leucomela, Michx.

Physcia leucomela, Michx., Fl. Bor.-Am., ii. p. 326; Nyl., Synop. Lich., i. p. 415, var. *angustifolia*; Mey. et Flot. in Nov. Act. Nat. Cur., xix., Suppl., i. p. 221; Crombie in Journ. Linn. Soc. Lond., xx. p. 83.

Borrera leucomela, Acharius, Lich. Univ., p. 499.

ASCENSION.—On the ground, amongst rocks—*Moseley.*

Widely spread.