

RAMALINEI.

Ramalina scopulorum, Acharius.

Ramalina scopulorum, Acharius; Nyl., Synop. Lich., i. p. 292; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 156, et in Melliss's St Hel., p. 374.

ST HELENA. *Melliss.*

From Lapland and Iceland to the Cape of Good Hope; also in North America, China, New Zealand, &c.

Ramalina polymorpha, Acharius.

Ramalina polymorpha, Acharius; Nyl., Synop. Lich., i. p. 293; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 156, et in Melliss's St Hel., p. 374.

ST HELENA. *Melliss.*

Widely diffused.

Ramalina pollinaria, Acharius.

Ramalina pollinaria, Acharius; Nyl., Synop. Lich., i. p. 296; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 156, et in Melliss's St Hel., p. 374.

ST HELENA. *Burchell.*

Widely diffused.

Ramalina ceruchis, De Notaris.

Ramalina ceruchis, De Notaris; Nyl., Synop. Lich., i. p. 289; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 156, et in Melliss's St Hel., p. 375.

ST HELENA. *Burchell.*

South America.

Ramalina angulosa, Laur.

Ramalina angulosa, Laur.; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 156, et in Melliss's St Hel., p. 374 (*formæ* 3).

Ramalina scopulorum, var. *implexa*, Nyl., Synop. Lich., i. p. 293.

ST HELENA. *Burchell.*

South Africa.

STICTEI.

Stictina crocata, Nyl.

Stictina crocata, Nyl., Synop. Lich., i. p. 338; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 156, et in Melliss's St Hel., p. 375.

ST HELENA. *Melliss.*

Very widely spread.