

Stictina tomentosa, Nyl.

Stictina tomentosa, Nyl, Synop. Lich., i. p. 343 ; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 156, et in Melliss's St Hel., p. 375.

ST HELENA. *Burchell*; *Melliss*.

Sticta aurata, Acharius.

Sticta aurata, Acharius ; Nyl, Synop. Lich., i. p. 361 ; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 156, et in Melliss's St Hel., p. 375.

ST HELENA. *Burchell*; *Melliss*.

Very widely spread.

Ricasolia herbacea, De Notaris.

Ricasolia herbacea, De Notaris ; Nyl, Synop. Lich., i. p. 369 ; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 156, et in Melliss's St Hel., p. 375.

ST HELENA. *Burchell*.

Specimen fragmentary, and uncertain as to the species.

Widely spread.

PARMELIEI.

Parmelia saxatilis, Acharius.

Parmelia saxatilis, Acharius ; Nyl, Synop. Lich., i. p. 388 ; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 156, et in Melliss's St Hel., p. 376.

ST HELENA. *Burchell*; *Melliss*.

Widely spread.

Parmelia perlata, Acharius.

Parmelia perlata, Acharius ; Nyl, Synop. Lich., i. p. 379 ; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 156, et in Melliss's St Hel., p. 375.

ST HELENA. *Melliss*.

Cosmopolitan, except in the colder regions.

Parmelia lævigata, Acharius.

Parmelia lævigata, Acharius ; Nyl, Synop. Lich., i. p. 384 ; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 156, et in Melliss's St Hel., p. 376.

ST HELENA. *Burchell*.

Cosmopolitan, except in the colder regions.