

Physcia speciosa, Fries, var. *hypoleuca*.

Physcia speciosa, Fries; Nyl., Synop. Lich., i. p. 416; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 156, et in Melliss's St Hel., p. 376 (var. *hypoleuca*).

ST HELENA. *Burchell*; *Melliss*.

This variety is widely dispersed, especially in warm regions.

Physcia astroidea, Fries.

Physcia astroidea, Fries; Nyl., Synop. Lich., i. p. 426; Leighton in Melliss's St. Hel., p. 376.

ST HELENA. *Melliss*.

Central and Southern Europe, and North Africa.

Physcia picta, Nyl.

Physcia picta, Nyl., Synop. Lich., i. p. 430; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 157, et in Melliss's St. Hel., p. 376.

ST HELENA.—*Melliss*.

Widely spread.

LECANOREI.

Pannaria rubiginosa, Del.

Pannaria rubiginosa, Del.; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 157, et in Melliss's St Hel., p. 376.

ST HELENA. *Burchell*; *Melliss*.

Very widely spread.

Erioderma unguigerum, Fée.

Erioderma unguigerum, Fée; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 157, et in Melliss's St Hel., p. 377.

ST HELENA. *Burchell*; *Melliss*.

Burchell's St Helena specimens of this lichen in Kew Herbarium are named *Erioderma polycarpum* in the handwriting of Leighton; and one collected by Sir J. D. Hooker is named *Erioderma unguigerum* by Nylander.

Cora pavonia, Web.

Cora pavonia, Web.; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 157, et in Melliss's St Hel., p. 377.

ST HELENA. *Burchell*; *Melliss*.

West Indies; South America.