

Lecidea enteroleuca, Acharius.

Lecidea enteroleuca, Acharius, Lich. Univ., p. 177, et Synop. Lich., p. 19; Leighton in Melliss's St Hel., p. 378.

ST HELENA. *Melliss.*

Europe, &c.

Opegrapha leightonii, Cromb.

Opegrapha leightonii, Crombie in Leight. Lich. Fl. Brit., p. 409; Leighton in Melliss's St Hel., p. 378.

ST HELENA. *Melliss.*

Europe, &c.

Lepraria ochracea, T. & B.

Lepraria ochracea, T. & B.; Leighton in Melliss's St Hel., p. 379.

ST HELENA. *Melliss.*

Europe, &c.

Lepraria flava, Acharius.

Lepraria flava, Acharius, Lich. Univ., p. 663, et Meth. Lich., p. 180; Leighton in Melliss's St Hel., p. 379.

ST HELENA. *Melliss.*

FUNGI.

HYMENOMYCETES.

Agaricus (Amanita) pantherinus, DC.

Agaricus (Amanita) pantherinus, DC., Fl. Fr., vi. p. 52; Fries, Hymen. Europ., p. 21; Berkl. in Melliss's St Hel., p. 379.

ST HELENA. *Melliss.*

Europe; North America.

Agaricus (Psalliota) campestris, Linn.

Agaricus (Psalliota) campestris, Linn., Sp. Pl., 2d ed., p. 1641; Fries, Hymen. Europ., p. 279; Berkl. in Melliss's St Hel., p. 379.

ST HELENA.—Abundantly on the grassy plains at Deadwood, Man and Horse, New Ground, &c.—*Melliss.*¹

North and south temperate regions; also from Ceylon, Java, Guiana, and Brazil.

¹ Melliss mentions that a larger species of *Agaricus* inhabits the rotting beds of haystacks; and a still larger one, about eight or ten inches in diameter, is occasionally met with on the barren plains between Longwood and Turk's Cap.