

**Agaricus (Hypholoma) fascicularis, Huds.**

*Agaricus (Hypholoma) fascicularis*, Huds., Fl. Angl., 2d ed., p. 615; Fries, Hymen. Europ. p. 291?,  
Berk. in Melliss's St Hel., p. 379.

ST HELENA. *Melliss.*

The true plant is widely dispersed in both north and south temperate regions.

**Schizophyllum commune, Fries.**

*Schizophyllum commune*, Fries, Syst. Mycol., i. p. 330; Berk. in Melliss's St Hel., p. 380.

ST HELENA. *Melliss.*

Ubiquitous.

**Polyporus lucidus, Fries.**

*Polyporus lucidus*, Fries, Hymen. Europ., p. 536; Berk. in Melliss's St Hel., p. 380.

ST HELENA.—Without locality—*Melliss.*

A cosmopolitan species.

**Polyporus sanguineus, Mey. ?**

*Polyporus sanguineus*, Mey., Esseq., p. 304; Fries, Syst. Mycol., i. p. 371?; Berk. in Melliss's St Hel.,  
p. 380.

ST HELENA. *Melliss.*

Specimen quite colourless from exposure to weather.

**Polyporus (Anodermei) induratus, Berk.**

*Polyporus (Anodermei) induratus*, Berk. in Melliss's St Hel., p. 380, t. 56.

ST HELENA.—Endemic. Abundant on the trunks of trees that lie rotting on the high  
land at 1600 to 2000 feet—*Melliss.*

“Pileo unguolato e tomentoso pallide sublaccato, poris elongatis albis minutis angulatis,  
dissepimentis tenuibus, contextu fibroso-sericeo.”

**Polyporus versicolor, Fries.**

*Polyporus versicolor*, Fries, Syst. Mycol., i. p. 368; Hymen., Europ., p. 568; Berk. in Melliss's St  
Hel., p. 380.

ST HELENA.—Without locality—*Melliss.*

This is a European species, and it is also recorded from India, Siberia, United States,  
Mexico, Cuba, Guadaloupe, Brazil, Juan Fernandez, Cape of Good Hope, New Zealand,  
Tasmania, Australia, and Java.