

RAMALINEI.

Ramalina yemenensis (Acharius), Nyl.

Ramalina yemenensis (Acharius), Nyl., Monogr. Ramal., p. 46; Crombie in Journ. Linn. Soc. Lond., xvi. p. 222.

INACCESSIBLE ISLAND. *Moseley.*

Arabia; India; Java; South Africa; South America; Australia; New Zealand.

Ramalina intermedia, Del.

Ramalina intermedia, Del. in Nyl. Monogr. Ramal., p. 68; Crombie in Journ. Linn. Soc. Lond., xvi. p. 222.

INACCESSIBLE ISLAND. *Moseley.*

Central and Western Europe; North America; Rodriguez.

PARMELIEI.

Parmelia perforata, Acharius.

Parmelia perforata, Acharius, Meth. Lich., p. 217; Nyl., Synop. Lich., i. p. 377; Crombie in Journ. Linn. Soc. Lond., xvi. p. 222.

TRISTAN DA CUNHA and INACCESSIBLE ISLANDS. *Moseley.*

Western Europe; America; South Africa; Polynesia; Australia.

Parmelia perlata, Acharius.

Parmelia perlata, Acharius, Meth., p. 216; Nyl., Synop. Lich., i. p. 379; Crombie in Journ. Linn. Soc. Lond., xvi. p. 222.

TRISTAN DA CUNHA. *Moseley.*

Parmelia revolutella, Nyl.

Parmelia revolutella, Nyl. in Journ. Linn. Soc. Lond., xvi. p. 222.

TRISTAN DA CUNHA.—Endemic. On trees? very sparingly seen—*Moseley.*

STICTEI.

Stictina intricata, var. *thouarsii*, Nyl.

Stictina intricata, var. *thouarsii*, Nyl., Synop. Lich., i. p. 335; Crombie in Journ. Linn. Soc. Lond., xvi. p. 222.

Sticta thouarsii, Del., Stict., n. 25, t. 8, fig. 9; Bory in Duperr., Voy. "Coquille," Bot. Crypt., p. 236.

TRISTAN DA CUNHA. *Thouars*; *Moseley.*

The species is widely diffused.