

ENUMERATION OF THE PLANTS.¹

PHANEROGAMÆ.

RANUNCULACEÆ.

Ranunculus crassipes,² Hook. f.

Ranunculus crassipes, Hook. f., Fl. Antarct., p. 224, t. 81, et in Phil. Trans. Roy. Soc. Lond., clxviii p. 17.

KERGUELEN ISLAND.—Christmas Harbour, Observatory, and Swain's Bay, Royal Sound.

Ranunculus trullifolius, Hook. f.

Ranunculus trullifolius, Hook. f., Fl. Antarct., p. 226, t. 82, A., et in Phil. Trans. Roy. Soc. Lond., clxviii p. 17, t. 1.

KERGUELEN ISLAND.—In streamlets and lakes, Royal Sound, Swain's Bay, Betsy Cove.

Ranunculus moseleyi, Hook. f.

Ranunculus moseleyi, Hook. f. in Phil. Trans. Roy. Soc. Lond., clxviii p. 18, t. 2, fig. 1.

KERGUELEN ISLAND.—In the lake at Christmas Harbour.

CRUCIFERÆ.

Pringlea antiscorbutica, R. Br.

Pringlea antiscorbutica, R. Br. ; Hook. f., Fl. Antarct., p. 238, t. 90, 91 ; Kidder in Bull. U.S. Nat. Mus., No. 321 ; Oliver in Journ. Linn. Soc. Lond., xiv. p. 389 ; Dyer in Proc. Linn. Soc. Lond. 1874, xxxiv. ; Hook. f., in Phil. Trans. Roy. Soc. Lond., clxviii p. 18, t. 2, fig. 3.

KERGUELEN ISLAND.—Throughout the island.

¹ Restricted to a list, with the necessary references, and the distribution within the island, extracted from Sir Joseph Hooker's paper on the botany of the island (*Phil. Trans. Roy. Soc. Lond.*, clxviii.), where the various collectors are enumerated. Particulars of the general distribution of the vascular plants will be found in a tabular form in the Report on the Botany of Heard Island (p. 244).

² Although the original specimens of this are very distinct from the ordinary *Ranunculus biternatus*, intermediate forms have been collected, and we have authority for saying that Sir Joseph Hooker would now give it the rank of a variety rather than an independent species.