

	Leach = <i>Terebratulina Grevillei</i> , T. Woods = <i>Terebratulina marginata</i> , Risso. = <i>Terebratulina quadrata</i> , Risso. = <i>Anomia retusa</i> , Lin. = <i>Delthyris spatula</i> , Menke.	Torridon, Scotland; off Belfast; off Finisterre and Croix de Gavie; Morbihan; Cape Breton, opposite southern extremity of Dep. des Landes, France; Adventure Bank, off Guetaria, Spain; N.E. coast of Jamaica; Korea; Australia. <i>Fossil</i> : Upper Tertiaries, Coralline Crag, Scotland; Scandinavia; Belgium; South Italy; Azores.
50 to 130	* <i>Terebratulina caput-serpentis</i> , var. <i>mediterranea</i> , Jeff., Proc. Zool. Soc., 1878. This seems to be a well-marked variety.	Mediterranean. <i>Fossil</i> : Sicily, South of Italy.
51 to 130	* <i>Terebratulina caput-serpentis</i> , var. <i>septentrionalis</i> , Couthouy, Boston Journ. Nat. Hist., vol. ii. p. 55, 1838. By some malacologists this is considered to be a distinct species, by others as only a variety of <i>T. caput-serpentis</i> . It has been several times well figured.	North-East Coast of America; off Halifax, Massachusetts Bay, Coast of Maine; East Port at low water, off Isle of Shoals; Grand Manan; off Cape of Good Hope; West of Kerguelen Island.
82	* <i>Terebratulina</i> (undetermined species), Dav., Proc. R. Soc., vol. xxvii., 1878.	Off Philippines (Challenger Expedition).
58 to 55	<i>Terebratulina japonica</i> , Sow., Thes. Conch., p. 344, pl. lxvi. figs. 7, 8, 1847.—(?) = <i>Ter. abyssicola</i> , Reeve, Voy. Samarang, p. 72, pl. xxi. fig. 5, 1850, (?) = <i>T. angulata</i> , Reeve, id., pl. xxi. fig. 2, 1850. Seems to be very variable in its shapes.	Japan; Gulf of Korea; China Seas.
26 to 63	(?) <i>Terebratulina cumingi</i> , Dav., Proc. Zool. Soc., p. 79, pl. xiv. figs. 17-19, 1852. Still an uncertain species, perhaps distinct, but thought by Mr Dall to be a variety or young <i>T. japonica</i> (?).	China Seas.
500	(?) <i>Terebratulina trigona</i> , Jeffreys, Proc. Zool. Soc., p. 402, pl. xxii. fig. 3, 1878. Requires more study, perhaps a young shell of some other species (?).	Coast of Portugal, dredged by Mr Kent in Dr Marshall Hall's yacht "Norma," in 1870.
340 to 430	(?) <i>Terebratulina tuberata</i> , Jeffreys, Proc. Zool. Soc., p. 401, pl. xxii. fig. 2, 1878.	Josephine Bank ("Josephine" Expedition and "Porcupine" Expedition, 1870).
5 to 50	* <i>Terebratulina</i> (?) <i>patagonica</i> , Gould, Proc. Boston Soc. Nat. Hist., 1850. The genus to which this species should be referred is still uncertain. I have never seen a specimen of it, it is stated to be striated, and looks as if it belonged to the sub-genus <i>Terebratulina</i> .	Orange Harbour, Patagonia.
70 to 471	<i>Terebratulina cailletii</i> , Crosse, Journ. Conchyl., p.	Florida; Tortugas, West In-